

On physical foundations and observational effects of cosmic rotation

Yuri N. Obukhov

Department of Theoretical Physics,

Moscow State University, 117234 Moscow, Russia

Abstract

An overview of the cosmological models with expansion, shear and rotation is presented. Problems of the rotating models are discussed, their general kinematic properties and dynamical realizations are described. A particular attention is paid to the possible observational effects which could give a definite answer to the question: is there evidence for the rotation of the universe?

1. INTRODUCTION

In his lecture “Spin in the Universe” [432], E.T. Whittaker has drawn attention to the following questions: *“Rotation is a universal phenomenon; the earth and all the other members of the solar system rotate on their axes, the satellites revolve round the planets, the planets revolve round the Sun, and the Sun himself is a member of the galaxy or Milky Way system which revolves in a very remarkable way. How did all these rotary motions come into being? What secures their permanence or brings about their modifications? And what part do they play in the system of the world?”*

The standard Friedman-Lemaître cosmological models are based on very idealized assumptions about the spacetime structure and matter source. The Friedman-Robertson-Walker metrics describe an exactly isotropic and homogeneous world filled with an ideal fluid (radiation and dust are the well known particular cases). Although these assumptions

are qualitatively supported by observations, it seems doubtful that the geometry of the universe and its physical contents are so finely tuned starting with the very Big Bang till the modern stage of the evolution. A reasonable step is to look for more general and realistic models with the assumptions of isotropy, homogeneity, and ideal fluid matter contents, relaxed. Here we will not discuss inhomogeneous cosmologies which are reviewed, e.g., in [238,239].

Already in the early years of the general relativity theory, attempts were made to construct (exact or approximate) solutions of the gravitational field equations with rotating matter sources (see, e.g. [225,11,12,231,407]). Lanczos [225] appears to be the first who considered the universe as the largest possible rotating physical system in a model of a rigidly rotating dust cylinder of an infinite radius. Dust density in this solution (which was later rederived by van Stockum [407]) diverges at radial infinity, which was a serious problem.

In 1946, Gamow [112] stressed a possible significance of the cosmic rotation for explaining the rotation of galaxies in the theories of galaxy formation, see also his discussion in [113]. Independently, Weizsäcker [430] initiated the study of the primordial turbulence and of its role in the formation of the universe's structure. Soon after this, K. Gödel [116] had proposed a stationary cosmological model with rotation in the form (coordinates $\{t, x, y, z\}$, $a = \text{const}$):

$$ds^2 = a^2 \left(dt^2 - 2e^x dt dy + \frac{1}{2} e^{2x} dy^2 - dx^2 - dz^2 \right). \quad (1.1)$$

In the Gödel's model, the matter is described as dust with the energy density ε , and the cosmological constant Λ is nontrivial and negative (i.e. its sign is opposite to that introduced by Einstein). The angular velocity ω of the *cosmic rotation* in (1.1) is given by $\omega^2 = \frac{1}{2a^2} = 4\pi G\varepsilon = -\Lambda$, with G as Newton's gravitational constant. For many years this model became a theoretical "laboratory" for the study of rotating cosmologies, see e.g. [6,429,27,79,152,153,217,336–338,406,72]. The stationary solution (1.1) is distinguished among other spatially homogeneous models: the so-called Gödel theorem (complete proof is given in [300]) states that the only spatially homogeneous models with zero expansion and

shear are Einstein's static universe and Gödel model (1.1).

Subsequently, Gödel [117] himself outlined a more physical *expanding* generalizations of (1.1), although without giving explicit solutions. Later a considerable number of exact and approximate models with rotation and with or without expansion were developed. At the end of this paper one can find a list of references which may be considered as a replacement of a detailed review of the previous work on rotating cosmology.

Here we would like to mention only briefly some earlier papers. Maitra [242] gave a generalization of van Stockum's solution which is cylindrically symmetric inhomogeneous stationary dust-filled world with rotation and shear. Maitra formulated a simple criterion of absence of the closed timelike curves. Wright [437] presented another inhomogeneous cylindrically symmetric solution for dust *plus* cosmological term.

Dust and cosmological constant represented the matter source in the series of papers by Schücking and Ozsváth [295–304], where the spatially homogeneous models with expansion, rotation and shear were considered. The closed Bianchi type IX solution ([299] with stationary limit [295,296]) has drawn a special attention. Most general anisotropic Bianchi type IX model was studied by Matzner [261,262] who took ideal fluid plus collisionless massless particles (photons and neutrinos) as matter source. The approximate solution was given. Later, Rebouças and de Lima [359] described the exact analytic Bianchi type IX solution with expansion, shear and rotation for a source represented by dust with heat flow and cosmological constant. Analogously, Sviestins [412] presented an exact Bianchi type IX solution for ideal fluid plus heat flow. In an interesting recent paper, Grøn [127] gave a *shear-free* Bianchi type IX solution in absence of matter (but with cosmological constant) which can be considered as a rotating generalization of de Sitter spacetime. In fact, the latter is recovered asymptotically after the quick decay of the vorticity.

The wide spectrum of spatially homogeneous models includes approximate and exact solutions for all Bianchi types and different matter contents (from simple dust to sometimes quite exotic energy-momentum currents). To mention but a few, Ruzmaikina-Ruzmaikin [375], Demiański-Grishchuk [88], and Batakis-Cohen [36] presented approximate solutions

of Bianchi type V and VII with expansion, shear and vorticity for the ideal fluid in non-comoving coordinates. Bradley and Sviestins [51] added heat flow to the ideal fluid source in their analysis of a Bianchi type VIII model. In a series of papers, Fennelly [104–108] has studied Bianchi models of different types for homogeneous universes filled with perfect charged (conducting) fluid and magnetic field. A neutral ideal fluid plus electromagnetic field and null radiation represented the matter source in a shear-free expanding and rotating solution of Patel-Vaidya [422] which can be interpreted as a standard Friedman-Robertson-Walker cosmology with rotating matter [130]. More sophisticated matter source may include viscous fluid and scalar fields, see, e.g., recent papers [246–256]. More exact solutions, which we will not discuss in detail, [7–9, 36, 51, 350, 283, 354, 356, 357, 370, 412, 85] were reported in the literature. Most of these models were overviewed by Krasinski [198–202], more recent review of generalized cosmologies with shear and rotation see in [206–208].

It is worthwhile to notice that along with the constant and deep interest in the rotating cosmologies, historical development has revealed several difficulties which were considered by the majority of relativists as the arguments against the models with nontrivial cosmic rotation. One of the problems was the stationarity of Gödel’s model (1.1). Apparent expansion of the universe is usually related to the fact of the red shift in the spectra of distant galaxies, and thus all the standard cosmological models are necessarily non-stationary. It was discovered, however, that it is impossible to combine pure rotation and expansion in a solution of the general relativity field equations for a simple physical matter source, such as a perfect fluid [95–97]. There are two ways to overcome that difficulty: one should either take a more general energy-momentum or to add cosmic shear, the explicit examples were already mentioned above. The possibility of combining cosmic rotation with expansion was the first successful step towards a realistic cosmology.

Another problem which was discovered already by Gödel himself, was related to causality: the spacetime (1.1) admits closed timelike curves. This was immediately recognized as an unphysical property because the causality may be violated in such a spacetime, see relevant discussion in [118, 143, 134, 135, 227, 336, 406]. Considerable efforts were thus focused

on deriving completely causal rotating cosmologies. In his last work devoted to rotation, Gödel without proof mentions the possibility of positive solution of the causality problem [117]. First explicit solutions were reported later [242,297]. Maitra [242] formulated a simple criterion for the (non)existence of closed timelike curves in rotating metrics which we will discuss below. Eventually, it became clear that the breakdown of causality is not inevitable consequence of the cosmic vorticity, and many of the physically interesting rotating models were demonstrated to be completely causal. That was the second important step in the development of the subject.

The discovery of the microwave background radiation (MBR) has revealed the remarkable fact that its temperature distribution is isotropic to a very high degree. This fact was for a long time considered as a serious argument for isotropic cosmological models and was used for obtaining estimates on the possible anisotropies which could take place on the early stages of the universe's evolution. In particular, homogeneous anisotropic rotating cosmologies were analyzed in [142,81,33,32,180,435], and very strong upper limits on the value of the cosmic rotation were reported. Unfortunately, usually in these studies the effects of vorticity were not separated properly from the cosmic shear.¹ For the first time, in [287,288], the effects of pure cosmic rotation were carefully separated from shear effects. The most significant result is the demonstration that in a large class of spatially homogeneous rotating models the pure rotation leads neither to an anisotropy of the MBR temperature nor to a parallax effects, although the geometry remains completely causal. A typical representative of that class is the so called Gödel type metric with rotation and expansion (4.1) which is considered below. Most recent analyzes, based also on the COBE data on the anisotropy of MBR, see [61,237,180], again estimated not a pure rotation but rather a shear.

The last but not least problem is the lack of direct observational evidence for the cosmic

¹Even now in the literature one can often encounter statements like: “If the universe had vorticity, the microwave background would be anisotropic...” [403].

rotation. Attempting at its experimental discovery one should study possible systematic irregularities in angular (ideally, over the whole celestial sphere) distributions of visible physical properties of sources located at cosmological distances. Unfortunately, although a lot of data is already potentially accumulated in various astrophysical catalogues, no convincing analysis was made in a search of the global cosmic rotation. Partially this can be explained by the insufficient theoretical study of observational cosmology with rotation. To our knowledge, till the recent time there were only few theoretical predictions concerning the possible manifestations of the cosmic rotation. Mainly these were the above mentioned estimates of MBR anisotropies [142,81,32,180] (also of the *X*-ray background anisotropies [435]) and the number counts analyses [225,117,431,265,103]. Besides that, it was pointed out in [417,373] that, in general, parallax effects can serve as a critical observational test for the cosmic rotation.

Very few purely empirical analyses (without constructing general relativistic models) of the angular distributions of astrophysical data are available which interpreted the observed systematic irregularities as the possible effects of rotation. Of the early studies, let us mention the reports of Mandzhos and Tel'nyuk-Adamchuk [243–245] on the preferential orientation of clusters of galaxies, Valdes et al [424] on the search of coherent ellipticity of galaxy images, and Andriasyan [16,17] on flattening of distant galaxies.

Birch [46,47,82] (Jodrell Bank Observatory) reported on the apparent anisotropy of the distribution of the observed angle between polarization vector and position of the major axis of radio sources, and related this global effect (using heuristic arguments) to the nontrivial cosmic rotation. Subsequently, these observations were disputed from a statistical theory viewpoint [335,43,44], but the analysis using indirectional statistics [173] confirmed Birch's results. It seems worthwhile to mention that the independently data from the Byurakan radio observatory [16,17] indicated the same anisotropy along, approximately, the same direction. These observations have stimulated the interest in rotating models; nevertheless, no further observations were reported until the most recently, when the new data appeared [279,279,347,348] claiming the same effect. Note however, that again the statistical signifi-

cance of observations is under discussion [428,94,229,68,234].

The theoretical analysis of the observational effects in the class of viable rotating cosmologies [287,187,288,194] has shown that the cosmic rotation produces a typical dipole anisotropy effect on the polarization of electromagnetic waves. Considering the case of the distant radio sources, one finds the dipole distribution [187,194] $\eta = \omega_0 r \cos \theta$, see (7.32) below, for the angle η between the vector of polarization and an observed direction of the major axis of the image of a radio source. Here ω_0 is the present value of the cosmic rotation, r is the (apparent area) distance to a source, and θ is the spherical angle between the rotation axis and the direction to the source, Z is the red shift. The data of Birch and the new data of Nodland and Ralston [279,279,347,348] may be extremely important for improving the direct estimates of vorticity.

The study of observational effects in rotating and expanding cosmological models is essentially based on the kinematical properties of spacetime. The next step is to investigate the *dynamical* aspects of cosmological models with rotation, in other words, to construct viable models as solutions of the gravitational field equations. The ultimate goal is to achieve a self-consistent description of the physical matter sources and to establish a complete scenario of a realistic evolution of the universe, including the most early stages of the big bang. Already Lemaître [230] and Whittaker [432] raised questions about the nature of a rotating “primeval atom” from which the universe is supposed to emerge. A related problem is the developments of matter irregularities which is relevant to the galaxy and large-scale structure formation. Gamov [112,113,13] and Weizsäcker [430] stressed the importance of the cosmic vorticity in the early universe. Silk [391,392] was the first to demonstrate instability of rotating models to axial density perturbations and stability to perturbations in the plane of rotation. The idea that primordial vortical motions of cosmological matter gave rise to the formation of large-scale structure underlied the related study of the so-called whirl theory (cosmological turbulence) developed some time ago by Ozernoy and Chernin [73–75,77,419,219–223,294], see also [276,293,22,83,84,138,139,393]. In the recent paper Li [232] has demonstrated that the global cosmic rotation can provide a natural origin of the

rotation of galaxies, and the empirical relation $J \sim M^{5/3}$ between the angular momentum J and the mass M of galaxies can be derived from the cosmological vorticity. [See Brosche [57–60] for empirical data, and Tassie [413] for the string-theoretical discussion].

Of the recent theoretical developments, it is worthwhile to mention the work [123] in which Grishchuk has demonstrated that the rotational cosmological perturbation can be naturally generated in the early universe through the quantum-mechanical “pumping” mechanism similar to the primordial gravitational waves. These perturbations were considered as the source of the large-angular-scale anisotropy of MBR, and comparison with observations was made. However, as was recognized by Grishchuk, the pumping mechanism only works when the primeval cosmological medium can sustain torque oscillations. No attempt was made to study the possible physical nature of the relevant cosmological matter. Exploiting the close relation between rotation and spin, one can try to attack this problem using the models of continua with microstructure. Self-consistent variational general relativistic theory of spin fluids was constructed in [289–291]. The investigation of the development of small perturbations for the models with spinning cosmological matter [291] yielded a generalization and refinement of the result by Silk [392] about instability to perturbations along the spin.

Most of the dynamical realizations of rotating cosmological models were obtained in the framework of general relativity theory as the exact or approximate solutions of Einstein’s gravitational field equations in four dimensions. However, we can mention also the study of higher-derivative extensions of general relativity [1–4,157], solutions in 5 dimensions [218,365], in 3 dimensions [133,433,434], and in string-motivated effective cosmology [30,31]. On the other hand, rotation, spin and torsion are closely interrelated in the Poincaré gauge theory of gravity [148,149,341,168,266], and hence it is quite natural to study cosmologies with rotation within the gauge gravity framework. Exact rotating solutions in Einstein-Cartan theory of gravity with spin and torsion were described, e.g., in [267,40,90,110,15,189,289,308,309,329,400,401]. General preliminary analysis of the separate

stages of the universe's evolution was made in our works [164–166], while in [189,167] the complete cosmological scenarios are considered.

Summarizing the present state of the problem, it is noteworthy once again to quote Whittaker [432]: *“It cannot be said, however, that any of the mathematical-physical theories that have been put forward to explain spin (rotation) in the universe has yet won complete and universal acceptance; but progress has been so rapid in recent years that it is reasonable to hope for a not long-delayed solution of this fundamental problem of cosmology”*.

2. STATIONARY COSMOLOGICAL MODEL OF GÖDEL TYPE

The Gödel metric (1.1) represents a particular case of a wider family of stationary cosmological models described by the interval

$$ds^2 = dt^2 - 2\sqrt{\sigma}e^{mx}dtdy - (dx^2 + ke^{2mx}dy^2 + dz^2), \quad (2.1)$$

where m, σ, k are constant parameters. Clearly, $\sigma > 0$, and for definiteness, we choose $m > 0$. The metric (2.1) is usually called the model with rotation of the Gödel type. Coordinate z gives the direction of the global rotation, the magnitude of which is constant:

$$\omega = \sqrt{\frac{1}{2}\omega_{\mu\nu}\omega^{\mu\nu}} = \frac{m}{2}\sqrt{\frac{\sigma}{k+\sigma}}. \quad (2.2)$$

As we see, vanishing of m or/and σ yields zero vorticity.

Let us describe the geometry of a Riemannian spacetime (2.1) in detail. The determinant of the metric is equal $(\det g) = -(k + \sigma)e^{2mx}$, and hence we assume that $(k + \sigma) > 0$ in order to have the Lorentzian signature.

Technically, all calculations are simplified greatly when exterior calculus is used. It is convenient to choose at any point of the spacetime (2.1) a local orthonormal (Lorentz) tetrad h_μ^a so that, as usual, $g_{\mu\nu} = h_\mu^a h_\nu^b \eta_{ab}$ with $\eta_{ab} = \text{diag}(+1, -1, -1, -1)$ the standard Minkowski metric. This choice is not unique, and we will use the gauge in which

$$h_0^{\hat{0}} = 1, \quad h_2^{\hat{0}} = -\sqrt{\sigma}e^{mx}, \quad h_1^{\hat{1}} = h_3^{\hat{1}} = 1, \quad h_2^{\hat{2}} = e^{mx}\sqrt{k+\sigma}. \quad (2.3)$$

Hereafter a caret denotes tetrad indices; Latin alphabet is used for the local Lorentz frames, $a, b, \dots = 0, 1, 2, 3$. The local Lorentz coframe one-form is defined, correspondingly, by

$$\vartheta^a := h_\mu^a dx^\mu. \quad (2.4)$$

The curvature two-form for (2.1) is computed straightforwardly:

$$R^{ab} = \omega^2 \vartheta^a \wedge \vartheta^b + \frac{k m^2}{k + \sigma} \left(\delta_1^a \delta_2^b - \delta_1^b \delta_2^a \right) \vartheta^{\hat{1}} \wedge \vartheta^{\hat{2}} \quad \text{for} \quad a, b = 0, 1, 2. \quad (2.5)$$

It is worthwhile to note that the spacetime (2.1) has a formal structure of a product of the curved *three-dimensional* manifold with the curvature (2.5) times flat one-dimensional space (z -axis), which is reflected in the range of indices in (2.5).

The only nontrivial components (with respect to the local Lorentz frame (2.3)) of the Weyl tensor are as follows:

$$C_{\hat{0}\hat{1}\hat{0}\hat{1}} = C_{\hat{0}\hat{2}\hat{0}\hat{2}} = -C_{\hat{2}\hat{3}\hat{2}\hat{3}} = -C_{\hat{3}\hat{1}\hat{3}\hat{1}} = \frac{1}{2} C_{\hat{1}\hat{2}\hat{1}\hat{2}} = -\frac{1}{2} C_{\hat{0}\hat{3}\hat{0}\hat{3}} = \frac{m^2}{6} \left(\frac{k}{k + \sigma} \right). \quad (2.6)$$

One thus verifies that the model (2.1) belongs to type D according to Petrov's classification.

The first dynamical realization of the model (2.1) was obtained by Gödel [116]. In general relativity theory, metric satisfies Einstein's field equations:

$$G_{ab} := R_{ab} - \frac{1}{2} g_{ab} R = \Lambda g_{ab} + \kappa T_{ab}. \quad (2.7)$$

Gödel [116] considered the simplest matter source: ideal dust with the energy-momentum tensor $T_{ab} = \rho u_a u_b$. Substituting the four-velocity of the co-moving matter $u^a = \delta_0^a$, and using the components of the Einstein tensor,

$$G_{\hat{0}\hat{0}} = -\omega^2 \left(1 + 4 \frac{k}{\sigma} \right), \quad G_{\hat{1}\hat{1}} = G_{\hat{2}\hat{2}} = \omega^2, \quad G_{\hat{3}\hat{3}} = \omega^2 \left(3 + 4 \frac{k}{\sigma} \right), \quad (2.8)$$

one finds the parameters of the famous Gödel solution [compare with (1.1)]:

$$\omega^2 = -\Lambda, \quad \kappa \rho = 2\omega^2, \quad \frac{k}{\sigma} = -\frac{1}{2}. \quad (2.9)$$

As we see, both the cosmological constant Λ and the parameter k must be *negative*.

Without confining ourselves to a particular dynamical realization, we should study the whole class of stationary models (2.1). There is a wide symmetry group for that metric. Three evident isometries are generated by the Killing vector fields

$$\xi_{(0)} = \frac{\partial}{\partial t}, \quad \xi_{(1)} = \frac{\partial}{\partial y}, \quad \xi_{(2)} = \frac{\partial}{\partial z}. \quad (2.10)$$

These are all mutually commuting. However, one can find two more Killing vectors. Most easily this can be done with the help of the suitable coordinate transformations.

Without touching z coordinate, let us transform the rest three

$$(t, x, y) \longrightarrow (\bar{\tau}, \bar{r}, \bar{\varphi})$$

as follows:

$$e^{mx} = e^{\bar{\varphi}} \cosh(m\bar{r}), \quad (2.11)$$

$$ye^{mx} = \frac{\sinh(m\bar{r})}{m\sqrt{k+\sigma}}, \quad (2.12)$$

$$\tan \left[m\sqrt{\frac{k+\sigma}{\sigma}}(t - \bar{\tau}) \right] = \sinh((m\bar{r})). \quad (2.13)$$

Straightforward calculation gives a new form of the metric (2.1):

$$ds^2 = d\bar{\tau}^2 - d\bar{r}^2 - \frac{[k \sinh(m\bar{r}) + \sigma]}{m^2(k + \sigma)} d\bar{\varphi}^2 - dz^2 + \frac{2}{m} \sqrt{\frac{\sigma}{k + \sigma}} \sinh(m\bar{r}) d\bar{\tau} d\bar{\varphi} \quad (2.14)$$

$$= \left(d\bar{\tau} + \frac{1}{m} \sqrt{\frac{\sigma}{k + \sigma}} \sinh(m\bar{r}) d\bar{\varphi} \right)^2 - \left(d\bar{r}^2 + \frac{\cosh^2(m\bar{r})}{m^2} d\bar{\varphi}^2 + dz^2 \right). \quad (2.15)$$

We thus immediately find the *fourth Killing* vector, $\partial_{\bar{\varphi}}$. In the original coordinates it reads:

$$\xi_{(3)} = \frac{\partial}{\partial \bar{\varphi}} = \frac{1}{m} \frac{\partial}{\partial x} - y \frac{\partial}{\partial y}. \quad (2.16)$$

Alternatively, let us consider another transformation

$$(t, x, y) \longrightarrow (\tau, r, \varphi)$$

given by the formulae:

$$e^{mx} = \cosh(mr) + \cos \varphi \sinh(mr), \quad (2.17)$$

$$ye^{mx} = \frac{\sin \varphi \sinh(mr)}{m\sqrt{k+\sigma}}, \quad (2.18)$$

$$\tan \left[\frac{m}{2} \sqrt{\frac{k+\sigma}{\sigma}} (t - \tau) \right] = \frac{\sin \varphi}{\cos \varphi + \coth \left(\frac{mr}{2} \right)}. \quad (2.19)$$

The metric (2.1) in the new coordinates reads:

$$ds^2 = d\tau^2 - dr^2 - \frac{4 \sinh^2 \left(\frac{mr}{2} \right) [k \cosh^2 \left(\frac{mr}{2} \right) + \sigma]}{m^2(k+\sigma)} d\varphi^2 - dz^2 \quad (2.20)$$

$$+ \frac{4}{m} \sqrt{\frac{\sigma}{k+\sigma}} \sinh^2 \left(\frac{mr}{2} \right) d\tau d\varphi$$

$$= \left(d\tau - \frac{2}{m} \sqrt{\frac{\sigma}{k+\sigma}} \sinh^2 \left(\frac{mr}{2} \right) d\varphi \right)^2 - \left(dr^2 + \frac{\sinh^2(mr)}{m^2} d\varphi^2 + dz^2 \right). \quad (2.21)$$

Evidently ∂_φ is also the Killing vector field. In the original coordinates it looks somewhat complicate and can be represented as a linear combination of three vectors:

$$\frac{\partial}{\partial \varphi} = \frac{1}{m} \sqrt{\frac{\sigma}{k+\sigma}} \xi_{(0)} + \frac{1}{2m\sqrt{k+\sigma}} \xi_{(1)} - \sqrt{k+\sigma} \xi_{(4)}, \quad (2.22)$$

where we denoted the *fifth Killing* vector:

$$\xi_{(4)} = \frac{\sqrt{\sigma} e^{-mx}}{m(k+\sigma)} \frac{\partial}{\partial t} + y \frac{\partial}{\partial x} + \frac{1}{2} \left[\frac{e^{-2mx}}{m(k+\sigma)} - m y^2 \right] \frac{\partial}{\partial y}. \quad (2.23)$$

Three of the five Killing vectors (2.10), (2.16), (2.23) satisfy commutation relations

$$[\xi_{(3)}, \xi_{(1)}] = \xi_{(1)}, \quad [\xi_{(3)}, \xi_{(4)}] = -\xi_{(4)}, \quad [\xi_{(1)}, \xi_{(4)}] = m\xi_{(3)}. \quad (2.24)$$

Thus $\xi_{(1)}, \xi_{(3)}, \xi_{(4)}$ form the closed subalgebra isomorphic to $so(2,1)$. Two Killing vectors $\xi_{(0)}, \xi_{(2)}$ span the center of the isometry algebra, commuting with all its elements.

In order to get some insight into possible causal problems, it is helpful to compute squares of the Killing vectors. Straightforwardly we obtain:

$$(\xi_{(0)} \cdot \xi_{(0)}) = 1, \quad (2.25)$$

$$(\xi_{(1)} \cdot \xi_{(1)}) = -k e^{2mx}, \quad (2.26)$$

$$(\xi_{(2)} \cdot \xi_{(2)}) = -1, \quad (2.27)$$

$$(\xi_{(3)} \cdot \xi_{(3)}) = - \left(k y^2 e^{2mx} + \frac{1}{m^2} \right), \quad (2.28)$$

$$(\xi_{(4)} \cdot \xi_{(4)}) = - \frac{k}{4} \left(e^{mx} m y^2 + \frac{e^{-mx}}{m(k+\sigma)} \right)^2, \quad (2.29)$$

The vector field $\xi_{(0)}$ is always timelike, its integral curves coincide with the lines of coordinate time. The cases $k > 0$, $k < 0$ and $k = 0$ are essentially different, and it is reasonable to consider them separately.

For *positive* values of k the four Killing vector fields $\xi_{(1)}, \dots, \xi_{(4)}$ are strictly spacelike; the first three of them provide spatial homogeneity of the $t = \text{const}$ hypersurfaces. We will demonstrate the complete causality of the models with $k > 0$ later. In particular, the closed timelike curves are absent in these geometries.

For *negative* k , the vectors $\xi_{(1)}$ and $\xi_{(4)}$ become timelike, whereas $\xi_{(3)}$ can be both spacelike and timelike, depending on the values of spatial coordinates x, y . One can immediately verify the existence of timelike closed curves, which was first noticed already by Gödel [116]. Indeed, consider the closed curve $\{t(\varphi), x(\varphi), y(\varphi)\}$, with $0 \leq \varphi \leq 2\pi$, given by the coordinate transformation (2.17)-(2.19) for $\tau = \text{const}, r = \text{const}, z = \text{const}$. From (2.20) we find that for large enough r , when $\cosh^2(\frac{mr}{2}) > |\frac{\sigma}{k}|$, such a curve is timelike. By construction, it is closed, and this may lead to the violation of causality. Although in [72] it was shown that such curves are not geodesics, one can in principle imagine a motion of a physical particle along these curves under the action of certain forces. It is worthwhile to note that the form of the metric is similar in two different coordinate systems, compare (2.15) and (2.21). However, for $k < 0$ the coordinate $\bar{\varphi}$ curve (with $\bar{\tau} = \text{const}, \bar{r} = \text{const}, z = \text{const}$) is also timelike for $\cosh^2(mr) > |\frac{\sigma}{k}|$, but it is *not closed*.

The *special case* $k = 0$ was studied in detail by Rebouças and Tiomno [360], see also [364] and more recently [368]. The Riemannian curvature is simplified greatly for $k = 0$. In particular, the Weyl tensor (2.6) vanishes, and thus four-dimensional metric is conformally flat. At the same time, (2.5) reduces to $R^{ab} = \omega^2 \vartheta^a \wedge \vartheta^b$, with $a, b = 0, 1, 2$. The last relation describes a three-dimensional manifold of constant curvature, i.e. (anti) de Sitter spacetime. This space has much wider isometry group than $k \neq 0$ case. In addition to the 5 Killing vectors (2.10), (2.16), (2.23), we easily find 2 more:

$$\xi_{(5)} = \sin(mt) \frac{\partial}{\partial t} - \cos(mt) \frac{\partial}{\partial x} + \frac{e^{-mx}}{\sqrt{\sigma}} \sin(mt) \frac{\partial}{\partial y}, \quad (2.30)$$

$$\xi_{(6)} = \cos(mt) \frac{\partial}{\partial t} + \sin(mt) \frac{\partial}{\partial x} + \frac{e^{-mx}}{\sqrt{\sigma}} \cos(mt) \frac{\partial}{\partial y}. \quad (2.31)$$

Together with ξ_0 these vector fields form the second closed subalgebra isomorphic to $so(2, 1)$:

$$[\xi_{(0)}, \xi_{(5)}] = m\xi_{(6)}, \quad [\xi_{(0)}, \xi_{(6)}] = -m\xi_{(5)}, \quad [\xi_{(5)}, \xi_{(6)}] = -m\xi_{(0)}. \quad (2.32)$$

The six Killing vectors $\xi_{(A)}$, $A = 1, 3, 4, 0, 5, 6$ with the commutation relations (2.24), (2.32) describe the complete $so(2, 2)$ algebra of isometries (algebra of the conformal group of a three-dimensional (anti) de Sitter spacetime) [368]. It is worthwhile to note that both new Killing vectors are spacelike:

$$(\xi_{(5)} \cdot \xi_{(5)}) = (\xi_{(6)} \cdot \xi_{(6)}) = -1, \quad (2.33)$$

Starting from the metric in the form (2.21) with $k = 0$ inserted, we can make a simple coordinate transformation

$$(\tau, r, \varphi) \longrightarrow (\tau, r, \theta),$$

redefining the angular coordinate,

$$\varphi = \frac{m}{2} (\theta - \tau), \quad (2.34)$$

which brings the metric to an explicitly (anti) de Sitter form:

$$ds^2 = \cosh^2\left(\frac{mr}{2}\right) d\tau^2 - dr^2 - \sinh^2\left(\frac{mr}{2}\right) d\theta^2 - dz^2. \quad (2.35)$$

Note that $\omega = \frac{m}{2}$ for $k = 0$.

3. CLASS OF SHEAR-FREE COSMOLOGICAL MODELS WITH ROTATION AND EXPANSION

In [287,187] we considered a wide class of viable cosmological models with expansion and rotation. Let us describe it here briefly. Denoting $x^0 = t$ as the cosmological time and $x^i, i = 1, 2, 3$ as three spatial coordinates, we write the space-time interval in the form

$$ds^2 = dt^2 - 2a n_i dx^i dt - a^2 \gamma_{ij} dx^i dx^j, \quad (3.1)$$

where $a = a(t)$ is the scale factor, and

$$n_i = \nu_a e_i^{(a)}, \quad \gamma_{ij} = \beta_{ab} e_i^{(a)} e_j^{(b)}. \quad (3.2)$$

Here $(a, b = 1, 2, 3)$ ν_a, β_{ab} are constant coefficients, while

$$e^{(a)} = e_i^{(a)}(x) dx^i \quad (3.3)$$

are the invariant 1-forms with respect to the action of a three-parameter group of motion which is admitted by the space-time (3.1). We assume that this group acts simply-transitively on the spatial ($t = \text{const}$) hypersurfaces. It is well known that there exist 9 types of such manifolds, classified according to the Killing vectors $\xi_{(a)}$ and their commutators $[\xi_{(a)}, \xi_{(b)}] = C^c_{ab} \xi_{(c)}$. Invariant forms (3.3) solve the Lie equations $\mathcal{L}_{\xi_{(b)}} e^{(a)} = 0$ for each Bianchi type, so that models (3.1) are spatially homogeneous.

We consider (t, x^i) as comoving coordinates for the cosmological matter, that is the four-vector of average velocity of material fluid is equal $u^\mu = \delta_0^\mu$. It is worthwhile to note that u^μ is not orthogonal to hypersurfaces of homogeneity. Such a models are usually called “tilted”. The kinematical characteristics of (3.1) are as follows: volume expansion is

$$\vartheta = 3 \frac{\dot{a}}{a}, \quad (3.4)$$

nontrivial components of vorticity tensor are

$$\omega_{ij} = -\frac{a}{2} \hat{C}^k_{ij} n_k, \quad i, j = 1, 2, 3, \quad (3.5)$$

and shear tensor is trivial,

$$\sigma_{\mu\nu} = 0. \quad (3.6)$$

Hereafter the dot ($\dot{}$) denotes derivative with respect to the cosmological time coordinate t . Tensor $\hat{C}^k_{ij} = e^k_{(a)}(\partial_i e_j^{(a)} - \partial_j e_i^{(a)})$ is the anholonomy object for the triad (3.3); for I-VII Bianchi types values of its components numerically coincide with the corresponding structure

constants C^a_{bc} . The list of explicit expressions for $\xi_{(a)}$, $e^{(a)}$, C^a_{bc} , $\hat{C}^k_{ij}$ for any Bianchi type is given in [287].

For completeness, let us mention that cosmological models (3.1) have nontrivial acceleration

$$u^\nu \nabla_\nu u_\mu = (0, \dot{a} n_i). \quad (3.7)$$

We choose the constant matrix β_{ab} in (3.2) to be *positive definite*. This important condition generalizes results of Maitra [242], and ensures the absence of closed time-like curves.

One can immediately see that space-times (3.1) admit, besides three Killing vector fields $\xi_{(a)}$, a nontrivial *conformal* Killing vector

$$\xi_{\text{conf}} = a \partial_t. \quad (3.8)$$

All models in the class (3.1) have a number of common remarkable properties [287,187]:

Causality: Space-time manifolds (3.1) are completely causal.

Indeed, following [242], let us suppose the spacetime (3.1) contains a closed curve $x^\mu(s)$ where s is some parameter, $0 \leq s \leq 1$, which is timelike. The latter means that at any point on the curve the length of the velocity vector is strictly positive,

$$g_{\mu\nu} \frac{dx^\mu}{ds} \frac{dx^\nu}{ds} > 0. \quad (3.9)$$

Since the curve is closed, $x^\mu(0) = x^\mu(1)$, there exists a value s_0 at which $\frac{dt}{ds} = 0$. At this point one has

$$g_{\mu\nu} \frac{dx^\mu}{ds} \frac{dx^\nu}{ds} \Big|_{s=s_0} = -a^2 \beta_{ab} e_i^{(a)} e_j^{(b)} \frac{dx^i}{ds} \frac{dx^j}{ds}, \quad (3.10)$$

which is negative when β_{ab} is positive definite. This contradicts (3.9) and hence we conclude that our assumption was wrong and there are no closed timelike curves.

For example, the stationary Gödel type metric (2.1) represents a particular case of (3.1) with $a = 1$ and

$$\nu_a = (0, \sqrt{\sigma}, 0), \quad \beta_{ab} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad e_i^{(a)} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{mx} & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (3.11)$$

It is clear that β_{ab} in (3.11) is positive definite for $k > 0$. Negative values of k , as we saw explicitly in Sec. 2, allow for the closed timelike curves. In particular, this is true for the original Gödel solution (2.9).

Isotropy of MBR: The Microwave Background Radiation (MBR) is totally isotropic in (3.1) for any moment of the cosmological time t .

In the geometric optics approximation, light propagates along null geodesics, i.e. curves $x^\mu(\lambda)$ with certain affine parameter λ and tangent vector $k^\mu = dx^\mu/d\lambda$ which satisfy

$$k^\nu \nabla_\nu k^\mu = 0, \quad k^\mu k_\mu = 0. \quad (3.12)$$

As it is well known, see e.g. [95,96], the red shift Z , which reflects the dependence of frequency on the relative motion of source and observer, is given by the simple formula

$$1 + Z = \frac{(k^\mu u_\mu)_S}{(k^\mu u_\mu)_P}, \quad (3.13)$$

where the subscripts “ S ” and “ P ” refer, respectively, to the spacetime points where a source emitted a ray and an observer detected it.

MBR is characterized by the black-body spectrum. One can show [95] that the temperature of the black-body radiation depends on the red shift,

$$T_P = \frac{T_S}{1 + Z} = T_S \frac{(k^\mu u_\mu)_P}{(k^\mu u_\mu)_S}. \quad (3.14)$$

In general, as was noticed, e.g., in [142,81,32,33,25], the observed temperature depends on the direction of observation through the ratio of $(k^\mu u_\mu)$ ’s. However, it is not the case for the class of shear-free models (3.1). Indeed, from (3.12) we easily prove that the scalar product of the wave vector with any Killing vector or conformal Killing vector

is constant along the ray. In particular, for the conformal Killing (3.8) we immediately obtain the first integral of the null-geodesics equations (3.12):

$$k_\mu \xi_{\text{conf}}^\mu = \text{const} \quad \implies \quad a(t_P)(u^\mu k_\mu)_P = a(t_S)(u^\mu k_\mu)_S, \quad (3.15)$$

where we used the evident proportionality of the Killing vector to the average four-velocity of matter,

$$\xi_{\text{conf}}^\mu = a(t) u^\mu. \quad (3.16)$$

Substituting (3.15) into (3.14), one finds

$$T_P = \frac{T_S}{1 + Z} = T_S \frac{a(t_S)}{a(t_P)}. \quad (3.17)$$

Thus, in the class of cosmological models (3.1), MBR is completely isotropic, exactly like in the standard cosmology. As a consequence, the earlier strong restrictions of the global rotation derived from the isotropy of MBR [142,81,32,33,25], are not applicable to the class of models under consideration. At the same time, the result obtained by no means contradicts the earlier observations, all of which were made for cosmologies *with shear*. Here we have trivial shear (3.6), and thus one clearly concludes that it is shear, not vorticity, which is the true source of anisotropy of MBR.

No parallaxes: Rotation (3.5) does not produce parallax effects.

In a very interesting paper [141], Hasse and Perlick introduced the generalized notion of parallax in cosmology. Let S_1 and S_2 be any two sources, seen by an observer P . Provided all the three objects do not have peculiar motion with respect to the matter with the average four-velocity u^μ , one says of a parallax effect when the angle between the rays coming from S_1 and S_2 to P , changes in time. Six equivalent conditions were formulated in [141] which are sufficient for the absence of parallax. One can easily check that all of them are satisfied for the class of models (3.1). For example, according to the P(3) condition (in notation of [141]), the velocity u^μ should be proportional to some

conformal Killing vector. Indeed, this is the case, (3.16). Alternatively, according to P(4) there should exist a function f such that

$$\frac{1}{2}\mathcal{L}_u g_{\mu\nu} = (u^\lambda \partial_\lambda f) g_{\mu\nu} - u_{(\mu} \partial_{\nu)} f.$$

One verifies straightforwardly that $f = \ln a$. According to P(5), u^μ is shear-free and the one-form

$$\rho := \left(u^\nu \nabla_\nu u_\mu - \frac{\vartheta}{3} u_\mu \right) dx^\mu$$

is closed, $d\rho = 0$. From (3.7) and (3.4), we find $\rho = 0$, hence P(5) is trivially fulfilled. Finally, according to P(6), there should exist a “red shift potential”. One can prove that again the latter if given by the function $f = \ln a$.

We thus conclude, that the models (3.1) are parallax-free, and consequently, the value of vorticity cannot be determined from parallax effects, contrary to the claims of [373,417,418].

Summarizing, cosmological models with rotation and expansion (3.1) solve the first three problems of cosmic rotation, and the most strong limits on the cosmic rotation, obtained earlier from the study of MBR [142,81,32] and of the parallaxes in rotating world [417,373], are not true for this class of cosmologies. This class of metrics is rich enough, as it contains all kinds of worlds: open and closed ones with different topologies.

4. GÖDEL TYPE EXPANDING COSMOLOGICAL MODEL

To the end of this paper we will consider now the natural non-stationary generalization of the original Gödel metric (1.1) which has drawn considerable attention in the literature. This generalized model is obtained from (2.1) by introducing the time-dependent scale factor $a(t)$,

$$ds^2 = dt^2 - 2\sqrt{\sigma}a(t)e^{mx}dtdy - a^2(t)(dx^2 + ke^{2mx}dy^2 + dz^2). \quad (4.1)$$

Here, $x^1 = x, x^2 = y, x^3 = z$, and the metric (4.1) evidently belongs to the family (3.1) with (3.11). As we saw already, the condition $k > 0$ guarantees the absence of closed time-like curves. The metric (4.1) is usually called the *Gödel type model* with rotation and expansion. Coordinate z gives the direction of the global rotation, the magnitude of which

$$\omega = \sqrt{\frac{1}{2}\omega_{\mu\nu}\omega^{\mu\nu}} = \frac{m}{2a}\sqrt{\frac{\sigma}{k+\sigma}} \quad (4.2)$$

decreases in expanding world [compare with (2.2)].

The three Killing vector fields are, recall (2.10) and (2.16),

$$\xi_{(1)} = \partial_y, \quad \xi_{(2)} = \partial_z, \quad \xi_{(3)} = \frac{1}{m}\partial_x - y\partial_y, \quad . \quad (4.3)$$

These satisfy commutation relations

$$[\xi_{(1)}, \xi_{(2)}] = \xi_{(2)}, \quad [\xi_{(1)}, \xi_{(3)}] = [\xi_{(2)}, \xi_{(3)}] = 0, \quad (4.4)$$

showing that the model (4.1) belongs to the Bianchi type III.

As before, it is convenient to choose at any point of the spacetime (4.1) a local orthonormal (Lorentz) tetrad h_μ^a . For simplicity, we will work with the immediate replacement of (2.3):

$$h_0^{\hat{0}} = 1, \quad h_2^{\hat{0}} = -a\sqrt{\sigma}e^{mx}, \quad h_1^{\hat{1}} = h_3^{\hat{3}} = a, \quad h_2^{\hat{2}} = ae^{mx}\sqrt{k+\sigma}. \quad (4.5)$$

From the point of view of the Petrov classification, one can verify that the Gödel type model is of the type *D*. Indeed, a straightforward calculation yields an obvious generalization of (2.6):

$$C_{\hat{0}\hat{1}\hat{0}\hat{1}} = C_{\hat{0}\hat{2}\hat{0}\hat{2}} = -C_{\hat{2}\hat{3}\hat{2}\hat{3}} = -C_{\hat{3}\hat{1}\hat{3}\hat{1}} = \frac{1}{2}C_{\hat{1}\hat{2}\hat{1}\hat{2}} = -\frac{1}{2}C_{\hat{0}\hat{3}\hat{0}\hat{3}} = \frac{m^2}{6a^2(t)}\left(\frac{k}{k+\sigma}\right). \quad (4.6)$$

5. DYNAMICAL REALIZATIONS

Several dynamical realizations (i.e. construction of exact cosmological solutions for the gravitational field equations) of the shear-free models (3.1), and in particular of the Gödel

type metric (4.1), are known. In the Einstein's general relativity theory the Gödel type models were described in [212,190,311–318,320], with different matter sources, whereas the Bianchi type II and III solutions (3.1) were obtained in [357,283] for the imperfect fluid with heat flow and cosmological constant.

In this section, we will describe a model [167] in which the gravitational field dynamics is determined by the minimal quadratic Poincaré gauge model. which is the closest extension of the Einstein's general relativity theory.

A. Field equations

In the framework of Poincaré gauge theory, gravitational field is described by the tetrad h_μ^a and the local Lorentz connection $\Gamma_{b\mu}^a$. In general case, gravitational Lagrangian is constructed as an invariant contraction from the curvature tensor

$$R^a{}_{b\mu\nu} = \partial_\mu \Gamma_{b\nu}^a - \partial_\nu \Gamma_{b\mu}^a + \Gamma_{c\mu}^a \Gamma_{b\nu}^c - \Gamma_{c\nu}^a \Gamma_{b\mu}^c, \quad (5.1)$$

and the torsion tensor

$$T^a{}_{\mu\nu} = \partial_\mu h_\nu^a - \partial_\nu h_\mu^a + \tilde{\Gamma}_{b\mu}^a h_\nu^b - \tilde{\Gamma}_{b\nu}^a h_\mu^b. \quad (5.2)$$

Note that here our notations for the connection, curvature and torsion are slightly different from [149]. In order to illustrate some common properties of cosmological models with rotation and expansion, here we consider the so called minimal quadratic Poincaré gauge gravity [167] which is the closest extension of Einstein's general relativity (we may also call such a model the generalized Einstein-Cartan theory). The corresponding Lagrangian reads:

$$L_g = -\frac{1}{16\pi G} \left[R + b R^2 + a_1 T^\alpha{}_{\mu\nu} T^\mu{}_{\alpha\nu} + a_2 T_{\alpha\mu\nu} T^{\mu\alpha\nu} + a_3 T_\mu T^\mu \right]. \quad (5.3)$$

Here $R = h_a^\mu h^{\nu b} R^a{}_{b\mu\nu}$ – the Riemann-Cartan curvature scalar, $T^\alpha{}_{\mu\nu} = h_a^\alpha T^a{}_{\mu\nu}$, and $T_\mu = T^\lambda{}_{\mu\lambda}$ – is the torsion trace, b, a_1, a_2, a_3 are the coupling constants.

Independent variation of the action (5.3) with respect to h_μ^a and $\tilde{\Gamma}_{b\mu}^a$ yields the field equations:

$$\begin{aligned}
& R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + 2bR \left(R_{\mu\nu} - \frac{1}{4}g_{\mu\nu}R \right) \\
& + a_1 \left(-\frac{1}{2}g_{\mu\nu}T_{\alpha\beta\gamma}T^{\alpha\beta\gamma} + 2T^\alpha_{\mu\beta}T_{\alpha\nu}{}^\beta - 2T_\alpha T_{\nu\mu}{}^\alpha - 2\nabla_\alpha T_{\nu\mu}{}^\alpha - T_{\mu\alpha\beta}T_\nu{}^{\alpha\beta} \right) \\
& + a_2 \left(-\frac{1}{2}g_{\mu\nu}T_{\alpha\beta\gamma}T^{\beta\alpha\gamma} + T^\alpha_{\beta\mu}T^\beta_{\alpha\nu} + T_\alpha T^\alpha_{\nu\mu} - T_\alpha T_{\mu\nu}{}^\alpha - \nabla_\alpha T_{\mu\nu}{}^\alpha + \nabla_\alpha T^\alpha_{\nu\mu} \right) \\
& + a_3 \left(\frac{1}{2}g_{\mu\nu}T_\alpha T^\alpha - \nabla_\nu T_\mu + g_{\mu\nu}\nabla_\alpha T^\alpha \right) = \kappa\Sigma_{\mu\nu},
\end{aligned} \tag{5.4}$$

$$\begin{aligned}
& - \left(\frac{1+2bR}{2} \right) (T^\alpha_{\mu\nu} + 2\delta^\alpha_{[\mu}T_{\nu]}) - 2b\delta^\alpha_{[\mu}\partial_{\nu]}R \\
& + (2a_1 - a_2)T_{[\mu\nu]}{}^\alpha - a_2T^\alpha_{\mu\nu} + a_3\delta^\alpha_{[\mu}T_{\nu]} = \kappa\tau^\alpha_{\mu\nu},
\end{aligned} \tag{5.5}$$

where $\kappa = 8\pi G$, with $\tau^\alpha_{\mu\nu}$ and $\Sigma_{\mu\nu}$ as the canonical tensors of spin and energy-momentum, respectively.

Equation (5.5) generalizes the well known algebraic relation between torsion and spin in the Einstein-Cartan theory. The tensors of torsion and spin may be decomposed into irreducible parts: trace T_μ and $\tau_\mu = \tau^\lambda_{\mu\lambda}$, pseudotrace $\overset{\vee}{T}_\mu = \frac{1}{2}\epsilon_{\mu\nu\alpha\beta}T^{\nu\alpha\beta}$ and $\overset{\vee}{\tau}_\mu = \frac{1}{2}\epsilon_{\mu\nu\alpha\beta}\tau^{\nu\alpha\beta}$, and trace-free parts $\bar{T}^\alpha_{\mu\nu}$ and $\bar{\tau}^\alpha_{\mu\nu}$:

$$T^\alpha_{\mu\nu} = \bar{T}^\alpha_{\mu\nu} + \frac{2}{3}T_{[\mu}\delta_{\nu]}^\alpha + \frac{1}{3}\epsilon^\alpha_{\mu\nu\lambda}\overset{\vee}{T}^\lambda, \tag{5.6}$$

$$\tau^\alpha_{\mu\nu} = \bar{\tau}^\alpha_{\mu\nu} + \frac{2}{3}\tau_{[\mu}\delta_{\nu]}^\alpha + \frac{1}{3}\epsilon^\alpha_{\mu\nu\lambda}\overset{\vee}{\tau}^\lambda. \tag{5.7}$$

Easy to see that eq. (5.5) decomposes into three equations for irreducible components:

$$(\mu_2 + 2bR)T_\mu + 3b\partial_\mu R = \kappa\tau_\mu, \tag{5.8}$$

$$(\mu_1 - bR)\overset{\vee}{T}_\mu = \kappa\overset{\vee}{\tau}_\mu, \tag{5.9}$$

$$(\mu_3 - bR)\bar{T}^\alpha_{\mu\nu} = \kappa\bar{\tau}^\alpha_{\mu\nu}, \tag{5.10}$$

where we denoted the following combinations of the coupling constants

$$\mu_1 = 2a_1 - 2a_2 - \frac{1}{2}, \tag{5.11}$$

$$\mu_2 = 1 - \frac{1}{2}(2a_1 + a_2 + 3a_3), \tag{5.12}$$

$$\mu_3 = -\left(\frac{1}{2} + a_1 + \frac{a_2}{2}\right), \tag{5.13}$$

which play important role in the determination of mass for quanta of torsion and couplings in all spin sectors of quadratic Poincaré gauge gravity model.

The eq. (5.8) shows that in contrast to the other irreducible parts, the torsion trace is characterized by the differential and not merely algebraic coupling. The Riemann-Cartan curvature scalar plays a role of its “potential”. Contraction of (5.4) yields an equation for R ,

$$-R + \frac{1}{2}(1 + 2\mu_3)\bar{T}_{\alpha\mu\nu}\bar{T}^{\alpha\mu\nu} + \frac{1}{6}(1 + 2\mu_1)\overset{\vee}{T}_\mu\overset{\vee}{T}^\mu + 2(1 - \mu_2)\left(\widetilde{\nabla}_\mu T^\mu - \frac{1}{3}T_\mu T^\mu\right) = \kappa\Sigma, \quad (5.14)$$

where $\Sigma = g_{\mu\nu}\Sigma^{\mu\nu}$ is the trace of the energy-momentum tensor. Inserting (5.8)-(5.10) into (5.14), we find a non-linear differential equation for the curvature scalar. As usual, hereafter we denote by tilde the purely Riemannian (torsion-free) geometrical objects and operators.

In view of specific nature of the torsion trace, it is convenient to separate its contribution to the eq. (5.4). Let us introduce

$$\hat{T}^\alpha{}_{\mu\nu} = \bar{T}^\alpha{}_{\mu\nu} + \frac{1}{3}\epsilon^\alpha{}_{\mu\nu\lambda}\overset{\vee}{T}^\lambda, \quad (5.15)$$

and hereafter hat “ $\hat{}$ ” will denote the corresponding geometrical quantities constructed with the help of the Riemann-Cartan connection with the trace-free torsion (5.15). The symmetric part of (5.4) then is rewritten as

$$\begin{aligned} & (1 + 2bR)\left(\hat{R}_{(\mu\nu)} - \frac{1}{2}g_{\mu\nu}\hat{R}\right) + \frac{1}{2}bR^2g_{\mu\nu} + \left(\frac{1 + 2\mu_3}{12}\right)T^\lambda\hat{T}_{(\mu\nu)\lambda} \\ & + a_1\left(-\frac{1}{2}g_{\mu\nu}\hat{T}_{\alpha\beta\gamma}\hat{T}^{\alpha\beta\gamma} + 2\hat{T}^\alpha{}_{\mu\beta}\hat{T}_{\alpha\nu}{}^\beta - \hat{T}_{\mu\alpha\beta}\hat{T}_\nu{}^{\alpha\beta} - 2\widehat{\nabla}_\alpha\hat{T}_{(\mu\nu)}{}^\alpha\right) \\ & + a_2\left(-\frac{1}{2}g_{\mu\nu}\hat{T}_{\alpha\beta\gamma}\hat{T}^{\beta\alpha\gamma} + \hat{T}_{\alpha\beta\mu}\hat{T}^{\beta\alpha}{}_\nu - \widehat{\nabla}_\alpha\hat{T}_{(\mu\nu)}{}^\alpha\right) \\ & + \left(\frac{8bR + 4\mu_2}{3}\right)\left(\nabla_{(\mu}T_{\nu)} - g_{\mu\nu}\nabla_\alpha T^\alpha - \frac{1}{6}g_{\mu\nu}T_\alpha T^\alpha\right) = \kappa\Sigma_{(\mu\nu)}, \end{aligned} \quad (5.16)$$

whereas the antisymmetric part of (5.4) is an identity in view of the geometric relation

$$2R_{[\mu\nu]} + (\nabla_\alpha + T_\alpha)\left(T^\alpha{}_{\mu\nu} + 2\delta_{[\mu}^\alpha T_{\nu]}\right) = 0, \quad (5.17)$$

and the angular momentum conservation law

$$(\nabla_\alpha + T_\alpha) \tau^\alpha_{\mu\nu} = \Sigma_{[\mu\nu]}. \quad (5.18)$$

Inserting the torsion $\overset{\vee}{T}_\mu$ and $\overline{T}^\alpha_{\mu\nu}$ from (5.9) and (5.10) into (5.15), and further into (5.16), one can write, similarly to the Einstein-Cartan theory, the effective Einstein equations for the metric. Spin of matter will contribute to the so called modified energy-momentum tensor.

B. Matter sources

We will describe cosmological matter by means of phenomenological model of a spinning fluid of Weyssenhoff and Raabe. This is a classical model of a continuous medium the elements of which are characterized, along with energy and momentum, by an intrinsic angular momentum (spin). Usually, hydrodynamical approach is considered as a good approximation to the description of realistic cosmological matter on the early, as well as on the later stages of universe's evolution. It seems natural to take into account spin (proper angular momentum) of elements of cosmological fluid, i.e., particles on early stage and galaxies and clusters of galaxies on later stages. In general case, an element of the Weyssenhoff fluid is characterized by the tensor of spin density $\tau_{\mu\nu}$, charge e and (proportional to spin) magnetic moment $\chi\tau_{\mu\nu}$ (χ – is a constant). The consistent variational theory of spin fluid in the Riemann-Cartan spacetime, developed in [289], yields the following currents in the right-hand sides of the gravitational field equations (5.4) and (5.5):

$$\begin{aligned} \Sigma_{\mu\nu} = & -pg_{\mu\nu} + u_\mu \left[u_\nu \left(\epsilon + p + \chi\tau_{\alpha\beta}F^{\alpha\beta} \right) + 2u^\alpha \nabla_\beta \tau^\beta_{\alpha\nu} \right] \\ & - F_{\mu\alpha} F_\nu{}^\alpha + \frac{1}{4}g_{\mu\nu} F_{\alpha\beta} F^{\alpha\beta} - 2\chi \left(\tau_{\mu\alpha} \tau_\nu{}^\alpha + \tau_{\nu\beta} F^{\alpha\beta} u_\alpha u_\mu \right), \end{aligned} \quad (5.19)$$

$$\tau^\alpha_{\mu\nu} = u^\alpha \tau_{\mu\nu}, \quad (5.20)$$

where u^μ is fluid's four-velocity, p is pressure, ϵ is the internal energy density, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the electromagnetic field created by and interacting with the electrodynamic characteristics of the medium.

Dynamics of the fluid is given by the rotational and translational equations of motion:

$$\begin{aligned}
(\nabla_\alpha + T_\alpha) \tau^\alpha_{\mu\nu} &= u_\mu u^\lambda \nabla_\alpha \tau^\alpha_{\lambda\nu} - u_\nu u^\lambda \nabla_\alpha \tau^\alpha_{\lambda\mu} \\
&\quad + \chi F^\alpha_\beta \left[(\delta^\beta_\mu - u^\beta u_\mu) \tau_{\alpha\nu} - (\delta^\beta_\nu - u^\beta u_\nu) \tau_{\alpha\mu} \right],
\end{aligned} \tag{5.21}$$

$$(\nabla_\nu + T_\nu) \Sigma^\nu_\mu - T^\alpha_{\mu\beta} \Sigma^\beta_\alpha - \tau^\nu_{\alpha\beta} R^{\alpha\beta}_{\mu\nu} = 0. \tag{5.22}$$

The self-consistent variational principle [289] yields also the Maxwell's equations for the electromagnetic field and the conservation law of number of particles:

$$\widetilde{\nabla}_\mu F^{\mu\nu} = e\rho u^\nu + 2\chi \widetilde{\nabla}_\mu \tau^{\mu\nu}, \tag{5.23}$$

$$\widetilde{\nabla}_\mu (\rho u^\mu) = 0, \tag{5.24}$$

where ρ is the particle density.

The system of differential and algebraic equations (5.4), (5.5), (5.19)-(5.24) is complete, and one can determine all the gravitational and material physical variables from this system. It is worthwhile to remind that spin satisfies the Frenkel condition,

$$u^\mu \tau_{\mu\nu} = 0. \tag{5.25}$$

The specific features of dynamics of the Weyssenhoff spin fluid in the Riemann-Cartan spacetime were studied in detail in [289].

C. Exact solution for the Gödel type universe

Let us describe cosmological model with rotation and expansion which provides an exact solution of the gravitational field equations. We will restrict ourselves to the shear-free class of spatially homogeneous metrics (3.1), and for definiteness, a particular case – the Gödel type model (4.1) – will be considered in detail.

The field equations (5.4) and (5.5) with the sources (5.19), (5.20) are most conveniently formulated with respect to (nonholonomic) local Lorentz frame. Let us choose the coframe (4.5). Its inverse reads:

$$h^0_0 = 1, \quad h^0_2 = \sqrt{\frac{\sigma}{k + \sigma}}, \quad h^3_3 = h^1_1 = \frac{1}{a}, \quad h^2_2 = \frac{1}{ae^{mx}\sqrt{k + \sigma}}, \tag{5.26}$$

Direct calculation of the local Lorentz connection $\tilde{\Gamma}^a_{b\mu} = h^a_\alpha h^\beta_b \tilde{\Gamma}^\alpha_{\beta\mu} + h^a_\alpha \partial_\mu h^\alpha_b$ (with $\tilde{\Gamma}^\alpha_{\beta\mu}$ as the Christoffel symbols) yields for the metric (4.1):

$$\tilde{\Gamma}^{\hat{0}}_{\hat{2}\hat{0}} = \tilde{\Gamma}^{\hat{1}}_{\hat{2}\hat{1}} = \tilde{\Gamma}^{\hat{3}}_{\hat{2}\hat{3}} = \frac{\dot{a}}{a} \sqrt{\frac{\sigma}{k + \sigma}}, \quad \tilde{\Gamma}^{\hat{1}}_{\hat{2}\hat{2}} = -\frac{m}{a}, \quad (5.27)$$

$$\tilde{\Gamma}^{\hat{2}}_{\hat{0}\hat{1}} = \tilde{\Gamma}^{\hat{2}}_{\hat{1}\hat{0}} = -\tilde{\Gamma}^{\hat{0}}_{\hat{1}\hat{2}} = \frac{m}{2a} \sqrt{\frac{\sigma}{k + \sigma}}, \quad (5.28)$$

where $\tilde{\Gamma}^a_{bc} = \tilde{\Gamma}^a_{b\mu} h^\mu_c$.

Since the direction of rotation (along the z axis) is apparently a distinguished one, it is natural to assume that the spin of the fluid and electromagnetic field are also oriented correspondingly along z . For the medium with vanishing magnetic moment of its elements ($\chi = 0$) one finds from (5.21)-(5.25):

$$\tau_{\hat{1}\hat{2}} = \tau_0 a^{-3}, \quad (5.29)$$

$$F_{\hat{1}\hat{2}} = B_0 a^{-2}, \quad (5.30)$$

$$\rho = \rho_0 a^{-3}, \quad (5.31)$$

where the integration constants are related by the consistency condition

$$m B_0 \sqrt{\frac{\sigma}{k + \sigma}} = e \rho_0. \quad (5.32)$$

We will now describe a solution of the gravitational field equations for which the Riemann-Cartan curvature scalar is assumed to be constant:

$$R = \text{const.} \quad (5.33)$$

Then in view of (5.25), eqs. (5.8)-(5.10) yield the irreducible parts of torsion:

$$T_\mu = 0, \quad \overset{\vee}{T}_\mu = \frac{\lambda_1}{2} \epsilon_{\mu\nu\alpha\beta} u^\nu \tau^{\alpha\beta}, \quad \bar{T}^\alpha_{\mu\nu} = \frac{\lambda_3}{3} (2u^\alpha \tau_{\mu\nu} + u_\mu \tau^\alpha_\nu - u_\nu \tau^\alpha_\mu), \quad (5.34)$$

where we denoted the constants

$$\lambda_1 = \frac{\kappa}{\mu_1 - bR}, \quad \lambda_3 = \frac{\kappa}{\mu_3 - bR}. \quad (5.35)$$

Nontrivial solution exists only when $\mu_1 \neq bR \neq \mu_3$, otherwise (5.9) and (5.10) lead to the vanishing spin. As for the eq. (5.8), one has two alternatives: if $\mu_2 + 2bR \neq 0$, then the torsion trace is zero, whereas if $\mu_2 + 2bR = 0$, then T_μ is arbitrary. However in both cases the torsion trace completely decouples from the eq. (5.16), hence without restricting generality we can put $T_\mu = 0$.

Using (5.34), we can rewrite (5.15) as

$$\hat{T}^\alpha_{\mu\nu} = \frac{1}{3} [(2\lambda_3 + \lambda_1) u^\alpha \tau_{\mu\nu} + (\lambda_3 - \lambda_1) u_\mu \tau^\alpha_\nu + (\lambda_1 - \lambda_3) u_\nu \tau^\alpha_\mu]. \quad (5.36)$$

Substituting this expression into (5.16), one finds

$$\begin{aligned} & a_1 \left(-\frac{1}{2} g_{\mu\nu} \hat{T}_{\alpha\beta\gamma} \hat{T}^{\alpha\beta\gamma} + 2\hat{T}^\alpha_{\mu\beta} \hat{T}^{\beta\alpha}_{\nu} - \hat{T}_{\mu\alpha\beta} \hat{T}^{\alpha\beta}_\nu - 2\hat{\nabla}_\alpha \hat{T}_{(\mu\nu)}^\alpha \right) \\ & + a_2 \left(-\frac{1}{2} g_{\mu\nu} \hat{T}_{\alpha\beta\gamma} \hat{T}^{\beta\alpha\gamma} + \hat{T}_{\alpha\beta\mu} \hat{T}^{\beta\alpha}_\nu - \hat{\nabla}_\alpha \hat{T}_{(\mu\nu)}^\alpha \right) = \\ & = \frac{1}{18} (2\lambda_3 + \lambda_1) [\lambda_1 (1 + 2\mu_1) - \lambda_3 (1 + 2\mu_3)] \tau_{\mu\alpha} \tau_\nu^\alpha \\ & + \left(\frac{\lambda_1 - 4\lambda_3}{36} \right) [\lambda_1 (1 + 2\mu_1) + 2\lambda_3 (1 + 2\mu_3)] u_\mu u_\nu \tau_{\alpha\beta} \tau^{\alpha\beta} \\ & + \frac{1}{24} [4\lambda_3^2 (1 + 2\mu_3) - \lambda_1^2 (1 + 2\mu_1)] g_{\mu\nu} \tau_{\alpha\beta} \tau^{\alpha\beta} + \lambda_3 (1 + 2\mu_3) \widetilde{\nabla}_\alpha [u_{(\mu} \tau_{\nu)}^\alpha]. \end{aligned} \quad (5.37)$$

After these computations, one finally finds that when the constant in (5.33) is equal $R = -1/2b$, the equations (5.16) are reduced to the algebraic relations between the pressure p , energy density ϵ , spin $\tau_{\alpha\beta}$ and Maxwell tensor $F_{\alpha\beta}$. From these equations, in addition to (5.32) we find a new relation between the integration constants,

$$B_0^2 = -m \tau_0 \sqrt{\frac{\sigma}{k + \sigma}}. \quad (5.38)$$

Without restricting generality, one can choose the parameter m to be positive, then the minus sign in (5.38) means that the spin and vorticity have the same direction. Thus $\tau_0 < 0$ [from (3.5), (4.5) and (5.26) we have $\omega_{\hat{1}\hat{2}} = -\frac{m}{2a} \sqrt{\frac{\sigma}{k+\sigma}}$]. In its turn, the direction of the magnetic field (sign of B_0) depends on the sign of the charge e .

With the help of (5.38) the equations (5.16) are reduced to the following two equations which describe the state of matter:

$$p(t) = -\frac{1}{8b\kappa} + \frac{B_0^2}{2a^4} + \left(\frac{\lambda_1 - 4\lambda_3}{6}\right) \frac{\tau_0^2}{a^6}, \quad (5.39)$$

$$\epsilon(t) = \frac{1}{8b\kappa} + \frac{3B_0^2}{2a^4} + \left(\frac{\lambda_1 - 4\lambda_3}{6}\right) \frac{\tau_0^2}{a^6}, \quad (5.40)$$

Finally, the evolution of the scale factor $a(t)$ is given by the equation (5.33). Substituting (4.5), (5.26), and (5.28) in it, one finds

$$\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} - \frac{m^2(3\sigma + 4k)}{12ka^2} - \left(\frac{k + \sigma}{36k}\right) (4\lambda_3^2 - \lambda_1^2) \frac{\tau_0^2}{a^6} - \frac{k + \sigma}{12kb} = 0. \quad (5.41)$$

As one can see, depending on the values of the coupling constants λ_3 and λ_1 , torsion can either accelerate or prevent the cosmological collapse. We will consider the latter possibility, assuming, for concreteness, that $4\lambda_3^2 \gg \lambda_1^2$.

Integrating (5.41), one gets

$$\dot{a}^2 = \frac{m^2(3\sigma + 4k)}{12k} + \frac{a^2(k + \sigma)}{24kb} - \frac{(k + \sigma)(4\lambda_3^2 - \lambda_1^2)\tau_0^2}{36ka^4} + \frac{\alpha}{a^2}. \quad (5.42)$$

The integration constant α is determined by the normalization of the scale factor $a = 1$ at the moment of a bounce ($t = 0$), when $\dot{a} = 0$,

$$\alpha = \left(\frac{k + \sigma}{36k}\right) (4\lambda_3^2 - \lambda_1^2) \tau_0^2 - \frac{m^2(3\sigma + 4k)}{12k} - \frac{k + \sigma}{24kb}. \quad (5.43)$$

D. Estimates of parameters of the cosmological model

In order to integrate the equation (5.42), one needs to know the constant parameters which enter that equation, because their numeric essentially affect the form of solution. All constants are uniquely determined by the data which describe modern state of the universe: the equation of state $p_t \approx 0$ (dust), energy density ϵ_t , Hubble constant $H_t (= \dot{a}/a)$ and the deceleration parameter $q_t (= -\ddot{a}a/\dot{a}^2)$ are estimated by

$$0.15 < \frac{\kappa\epsilon_t}{H_t^2} < 12,$$

$$4 \cdot 10^{-11} \text{ yr}^{-1} < H_t < 10^{-10} \text{ yr}^{-1},$$

$$0.01 < q_t < 1.$$

(Hereafter the index t denotes modern values of the physical quantities).

Previously, [46], the angular velocity was estimated as relatively small $\omega_t/H_t \approx 10^{-3}$. New analyses of observational data suggests higher estimates for the global vorticity, more close to H_t . For the qualitative analysis of the model let us choose

$$q_t = 0.01, \quad \frac{\kappa \epsilon_t}{H_t^2} = 0.18, \quad \frac{\omega_t}{H_t} = 0.1. \quad (5.44)$$

Besides the estimates (5.44), one needs also the values of the spin-torsion coupling constants. There is a large ambiguity which affects the scale of the space at the moment of a bounce. If one assumes, cf. Trautman [416], that the cosmological collapse stops on the scale of order of 1 cm, then we get from (5.44) an estimate

$$\frac{\kappa}{\lambda_1 - 4\lambda_3} \approx 2.3 \cdot 10^{27}. \quad (5.45)$$

Comparison of the equations (5.41)-(5.43) with (5.44)-(5.45) yields the modern value of the spin density

$$3 \cdot 10^8 \text{ g cm}^{-1} \text{ s}^{-1} < \tau_t < 7.6 \cdot 10^8 \text{ g cm}^{-1} \text{ s}^{-1}, \quad (5.46)$$

the geometric parameters of the model

$$\frac{k}{\sigma} \approx 71, \quad (5.47)$$

the coupling constant

$$b^{-1} = 0.36 H_t^2, \quad (5.48)$$

and the magnetic field

$$10^{-6} \text{ G} < B_t < 2.6 \cdot 10^{-6} \text{ G}. \quad (5.49)$$

The final integration of the equation (5.42) determines the law of evolution of the scale factor:

$$4\sqrt{\alpha_3} H_t t = (\text{const}) + \ln Z(a^2) + \frac{2\mu}{\widehat{k}\sqrt{-B_1 B_2}} \left[\frac{\gamma_1}{\gamma_1 + \gamma_2} F(\varphi; \widehat{m}) - C_1 \Pi(\varphi; n, \widehat{m}) \right], \quad (5.50)$$

where the integration constant (const) is such that $a = 1$ for $t = 0$, and we denote the function of the scale factor

$$Z(a^2) = \frac{(a^2 + \gamma_1)^2 + \sqrt{a^2(a^2 - 1)(a^2 - u_1)(a^2 - u_2)}}{(\gamma_1 + \gamma_2)(\gamma_1 - \gamma_2 + 2a^2)} - \frac{C_1 + C_2}{2}.$$

In eq. (5.50), the constant parameters are determined by:

$$\alpha_1 = \frac{m^2(3\sigma + 4k)}{12kH_t^2}, \quad \alpha_2 = \left(\frac{k + \sigma}{k}\right) \left(\frac{4\lambda_3^2 - \lambda_1^2}{36}\right) \frac{\tau_0^2}{H_t^2}, \quad \alpha_3 = \frac{k + \sigma}{12bkH_t^2};$$

$u_1 > u_2$ are the roots of the quadratic equation

$$u^2 + \left(1 + \frac{\alpha_1}{\alpha_3}\right)u + \frac{\alpha_2}{\alpha_3} = 0,$$

from which one constructs

$$\begin{aligned} \lambda_{\pm} &= -u_1 u_2 \left(\sqrt{1 - 1/u_1} \pm \sqrt{1 - 1/u_2} \right)^2, \\ B_1 &= \frac{\lambda_+ (1 - \lambda_-)}{\lambda_+ - \lambda_-}, \quad C_1 = \frac{\lambda_- (\lambda_+ - 1)}{\lambda_+ - \lambda_-}, \quad B_2 = \frac{1 - \lambda_-}{\lambda_+ - \lambda_-}, \quad C_2 = \frac{\lambda_+ - 1}{\lambda_+ - \lambda_-}, \\ \gamma_1 &= \sqrt{\frac{u_1 u_2}{1 - \lambda_-}}, \quad \gamma_2 = \sqrt{\frac{u_1 u_2}{1 - \lambda_+}}, \quad \mu^2 = -\frac{B_1}{C_1}, \quad \hat{k}^2 = \frac{\lambda_+}{\lambda_-}, \\ \widehat{m}^2 &= (\lambda_+ - \lambda_-) / \lambda_+, \quad n = B_1 \widehat{m}^2 = 1 - \lambda_-. \end{aligned}$$

Finally, $F(\varphi; \widehat{m})$ and $\Pi(\varphi; n, \widehat{m})$ are elliptic integrals of the first and the third kinds, respectively, with their argument given by

$$(1 - \hat{k}^{-2}) \sin^2 \varphi = 1 - \mu^{-2} \left(\frac{a^2 - \gamma_2}{a^2 - \gamma_1} \right)^2.$$

The complete picture of the cosmological evolution is obtained when one substitutes (5.44)-(5.49) into the above equations. In particular, one can verify that the age of the universe [estimated as the time since $t = 0$ till the modern stage characterized by (5.44)] is equal $T \approx H_t^{-1}$, in a good agreement with observational data.

The analysis of the evolution of the scale factor $a(t)$, see (5.50) and (5.42), shows that one can naturally distinguish several qualitatively different stages in the history of the universe. First stage is the shortest one and it describes a bounce in the vicinity of $t = 0$. There is no

initial singularity due to the dominating spin contribution in (5.42). At this stage, the fluid source is characterized by the approximate equation of state of stiff matter

$$p = \epsilon \approx \left(\frac{\lambda_1 - 4\lambda_3}{6} \right) \frac{\tau_0^2}{a^6}$$

The applicability of the classical (non-quantum) gravitational theory is guaranteed by the condition on the curvature invariant $\tilde{R}_{\alpha\beta\mu\nu}\tilde{R}^{\alpha\beta\mu\nu} \ll l_{\text{pl}}^{-4}$, which is satisfied at the moment of the bounce. The duration of that stage is quite small ($\ll 1$ s), since the spin term quickly decreases in (5.42) with the growth of the scale factor.

Next comes the second stage when the scale factor increases like $\sqrt{t}$, while the equation of state is of the radiation type, $p \approx \epsilon/3$. This “hot universe” expansion lasts until the size of the metagalaxy approaches $\approx 10^{27}$ cm. After this the “modern” stage starts with the effectively dust equation of state $p_t \approx 0$, $\epsilon_t \approx (2b\kappa)^{-1}$. Scale factor still increases, but the deceleration of expansion takes place. The final stage depends on the value of the cosmological term, and either the future evolution enters the eternal de Sitter type expansion, or expansion ends and a contraction phase starts.

Our purpose in this section was to demonstrate the exact solution which avoids the principal difficulties of old cosmological models with rotation and expansion. Certainly, one cannot claim that the results obtained describe the real universe, and consequently we have limited ourselves to a test type computations for the physical and geometrical parameters, without trying to find the best estimates. The main difficulty of this particular dynamical realization, in our opinion, is presented by the magnitude of magnetic field (5.49) which at the “modern” stage should be close to the upper limits established for the global magnetic field from astrophysical observations, see e.g. [24].

It seems worthwhile to note that (5.46) gives a rather big value for the spin density ($\approx 10^{35}\hbar/\text{cm}^3$). This is in a good agreement with the macroscopic interpretation of the cosmological spin fluid, the elements of which are spinning galaxies. Indeed, $\tau_t/\rho_t \approx 10^{107}\hbar$ is close to the observed magnitude of the angular momentum of a typical galaxy. Moreover, for the total spin of the metagalaxy occupying the volume $V \approx 10^{84}\text{cm}^3$, one finds $\tau_{\text{tot}} =$

$\tau_t V \approx 10^{120} \hbar$. This once again draws attention to the Large Number hypothesis of Dirac-Eddington and related questions, see [57–60,63,440] and [271–274].

6. NULL GEODESICS IN THE GÖDEL TYPE MODEL

Practically all the information about the structure of the universe and about the properties of astrophysical objects is obtained by an observer in the form of different kinds of electromagnetic radiation. Thus, in order to be able to make theoretical predictions and compare them with observations, it is necessary to know the structure of null geodesics in the cosmological model with rotation. All the models from the class (3.1) have three Killing vectors and one conformal Killing vector field. Hence the null geodesics equations (3.12) have four first integrals,

$$q_0 = \xi_{\text{conf}}^\mu k_\mu, \quad q_a = -\xi_{(a)}^\mu k_\mu, \quad a = 1, 2, 3. \quad (6.1)$$

Solving (6.1) with respect to k^μ , one obtains a system of ordinary first order nonlinear equations which can be straightforwardly integrated. Complete solution of the null geodesics equations in the Gödel type model is given in [186,287], and here we present only short description of null geodesics in (4.1).

To begin with, let us define convenient parametrization of null geodesics. Without losing generality (using the spatial homogeneity) we assume that an observer is located at the space-time point $P = (t = t_0, x = 0, y = 0, z = 0)$. Now, arbitrary geodesics which passes through P is naturally determined by its initial direction in the local Lorentz frame of observer at this point. In the tetrad (4.5) we may put

$$k_P^a = (h_\mu^a k^\mu)_P = (1, \sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta), \quad (6.2)$$

where θ, ϕ are standard spherical angles parametrizing the celestial sphere of an observer. Then from (3.8), (4.3), (6.2) and (6.1) one finds the values of the integration constants

$$q_0 = a_0,$$

$$q_1 = a_0(\sqrt{\sigma} + \sqrt{k + \sigma} \sin \theta \sin \phi), \quad (6.3)$$

$$q_2 = a_0 \cos \theta,$$

$$q_3 = \frac{a_0}{m} \sin \theta \cos \phi. \quad (6.4)$$

Generic null geodesics, initial directions of which satisfy $q_1/a_0 = (\sin \theta \sin \phi + \sqrt{\frac{\sigma}{k+\sigma}}) \neq 0$, are then described by

$$e^{-mx} = \frac{\sqrt{\sigma} + \sqrt{k + \sigma} \sin \theta \sin \Phi}{\sqrt{\sigma} + \sqrt{k + \sigma} \sin \theta \sin \phi}, \quad (6.5)$$

$$y = \frac{\sin \theta (\cos \Phi - \cos \phi)}{m(\sqrt{\sigma} + \sqrt{k + \sigma} \sin \theta \sin \phi)}, \quad (6.6)$$

$$z = \left(\frac{k + \sigma}{k} \right) \cos \theta \left[\int_{t_0}^t \frac{dt'}{a(t')} + \sqrt{\frac{\sigma}{k + \sigma}} \left(\frac{\Phi - \phi}{m} \right) \right], \quad (6.7)$$

where the function $\Phi(t)$ satisfies the differential equation

$$\frac{d\Phi}{dt} = -\frac{m}{a} \left(\frac{\sqrt{\frac{\sigma}{k+\sigma}} + \sin \theta \sin \Phi}{1 + \sqrt{\frac{\sigma}{k+\sigma}} \sin \theta \sin \Phi} \right) \quad (6.8)$$

with initial condition $\Phi(t_0) = \phi$.

For a detailed discussion of rays which lie on the initial cone $(\sin \theta \sin \phi + \sqrt{\frac{\sigma}{k+\sigma}}) = 0$ and different subcases of (6.5)-(6.8) see [287].

7. OBSERVATIONS IN ROTATING COSMOLOGIES

The qualitative picture of specific rotational effects which could be observed in the Gödel type model (4.1) is in fact independent of the dynamical behaviour of the scale factor $a(t)$. To some extent the same is true also for the quantitative estimates, especially if one uses the Kristian-Sachs formalism [214,240] in which all the physical and geometrical observable quantities are expressed in terms of power series in the affine parameter s or the red shift Z . In this case the description of observable effects on not too large (although cosmological) scales involves only the modern values (i.e. calculated at the moment of observation $t = t_0$) of the scale factor $a_0 = a(t_0)$, Hubble parameter $H_0 = (\dot{a}/a)_P$, rotation value $\omega_0 = \omega(t_0)$, deceleration parameter $q_0 = -(\ddot{a}a^2/\dot{a}^2)_P$, etc.

In this section we will outline possible observational manifestations of the cosmic rotation in the Gödel type universe. Estimates for the value of vorticity and for the direction of rotation axis can be found from the recent astrophysical data, see Sects. 7C and 7D.

A. Kristian-Sachs expansions in observational cosmology

Kristian and Sachs in the fundamental paper [214] have developed universal technique for the analysis of the observational effects in an arbitrary cosmological model. Essentially, their method is based on the theory of null geodesic (ray) bundles in Riemannian spacetime. The main advantage of the Kristian-Sachs approach is in its generality which makes it possible to consider cosmological tests even without the knowledge of explicit solution of the gravitational field equations. This is particularly convenient when one is interested in revealing additional (as compared to the standard Hubble expansion) geometrical and physical properties of the universe. In this section we will apply the method of Kristian and Sachs to the rotating cosmological models.

Let k^μ be a wave vector field which satisfies the null geodesics equation (3.12) in a Riemannian manifold. (The connection is torsion-free, and we will omit the tilde to simplify the notation). The starting point is the equation

$$\ddot{\Delta}^\mu = -R^\mu{}_{\alpha\nu\beta} \Delta^\nu k^\alpha k^\beta, \quad (7.1)$$

which determines the deviation vector $\Delta^\mu := \delta x^\mu$ of the rays in a bundle. Hereafter dots denote covariant derivatives along a ray: $(\dot{}) = k^\mu \nabla_\mu$, e.g., $\dot{\Delta}^\mu = D\Delta^\mu/ds = k^\nu \nabla_\nu \Delta^\mu$, etc.

The crucial idea of Kristian and Sachs is to find a solution of (7.1) in the form of a series in the affine parameter s . Correspondingly, all the physical and geometrical quantities are ultimately represented by series in s . Let us give some basic results of that approach, without going into calculational details.

We will denote the spacetime points of an observer and a source by P and S , respectively. Let us start with the red shift defined by (3.13). A simple Taylor expansion of the right-hand side yields:

$$Z = -\bar{s} \frac{(1)}{u} + \frac{\bar{s}^2}{2} \frac{(2)}{u} - \frac{\bar{s}^3}{6} \frac{(3)}{u} + \dots \quad (7.2)$$

Here we denoted:

$$\bar{s} := s (k^\mu u_\mu)_P, \quad K^\mu = \left(\frac{k^\mu}{k^\nu u_\nu} \right), \quad \frac{(n)}{u} := \left(K^{\mu_1} \dots K^{\mu_{n-1}} K^{\mu_n} \nabla_{\mu_1} \dots \nabla_{\mu_{n-1}} u_{\mu_n} \right)_P.$$

Note that the value s describes a position of the observer on a null geodesic connecting S and P , so that $s = 0$ corresponds to S . The minus signs in (7.2) arise when we expand $(u^\mu k_\mu)_S$ at P .

Let us consider a bundle of rays emitted from a point source S . We are interested in $\Delta^\mu(s)$ along a geodesic from S to P . One has $\Delta^\mu(0)$, and hence from (7.1), $\ddot{\Delta}^\mu(0) = 0$. The vector $\dot{\Delta}^\mu(0)$ is chosen to be orthogonal to k^μ and u^μ at S , i.e. we are studying the cross-section of the bundle which is orthogonal to the central ray in the rest frame of the source. Note that such an initial condition provides orthogonality $k^\mu \Delta_\mu = 0$ at any value of s , in view of (7.1).

The *area distance* is one of the central notions in the observational cosmology. If an observer P is viewing the distant source S with an intrinsic perpendicular area dA_S which subtends the solid angle $d\Omega_P$ at P , the area distance r between P and S is defined by

$$dA_S = r^2 d\Omega_P, \quad (7.3)$$

Another (reciprocal) area distance r_S between S and P is defined by $dA_P = r_S^2 d\Omega_S$, where the solid angle $d\Omega_S$ subtends the area dA_P at P . The *reciprocity theorem* [95,96,240] tells that these distances are related by

$$r_S^2 = r^2 (1 + Z)^2. \quad (7.4)$$

One can obtain r_S with the help of $\Delta^\mu(s)$ which describes the behaviour of the bundle of rays.

The deviation equation (7.1) determines iteratively the third and higher covariant derivatives of Δ^μ at S , and thus one construct the solution in the form of the Taylor series

$$\Delta^\mu(s) = s \dot{\Delta}^\mu(0) + \frac{s^3}{6} \ddot{\Delta}^\mu(0) + \dots$$

In order to find r_S , one needs actually only the length of Δ^μ . Straightforward calculation yields:

$$\begin{aligned} \Delta^\mu(s)\Delta_\mu(s) &= s^2 \dot{\Delta}^\mu(0)\dot{\Delta}_\mu(0) \left[1 - \frac{s^2}{6} b - \frac{s^3}{12} \dot{b} + O(s^4) \right]_S \\ &\quad + s^2 \dot{\Delta}^\mu(0)\dot{\Delta}^\nu(0) \left[-\frac{s^2}{3} c_{\mu\nu} - \frac{s^3}{6} \dot{c}_{\mu\nu} + O(s^4) \right]_S. \end{aligned} \quad (7.5)$$

The spacetime curvature contributes via the expressions

$$b := R_{\mu\nu} k^\mu k^\nu, \quad c_{\alpha\beta} := C_{\alpha\mu\beta\nu} k^\mu k^\nu, \quad (7.6)$$

higher order corrections are contained in the terms $O(s^4)$.

After some algebra, one finds from (7.5):

$$r_S^2 = s^2 (k^\mu u_\mu)_S \left[1 - \frac{s^2}{6} b - \frac{s^3}{12} \dot{b} + O(s^4) \right]_S. \quad (7.7)$$

Combining (3.13), (7.4), and (7.7), one can easily derive the series which represents the area distance r in terms of s .

However, since the value of the affine parameter s for the source S and the observer P cannot be measured directly, the series in powers of s are not particularly useful. Instead, Kristian and Sachs proposed to invert the relation (7.2), and then substitute $s = s(Z)$ in all other quantities. Direct calculation yields the following final series representation:

$$1 + Z = 1 - r \frac{(1)}{u} + \frac{r^2}{2} \frac{(2)}{u} - \frac{r^3}{6} \left(\frac{(3)}{u} + \frac{1}{2} B \right) + \dots, \quad (7.8)$$

where, replacing (7.6), $B := (R_{\mu\nu} K^\mu K^\nu)_P$. Inverting (7.8), one finds

$$r = \frac{Z}{-\frac{(1)}{u}} \left[1 - \frac{Z}{2} \frac{\frac{(2)}{u}}{\frac{(1)}{u^2}} + \frac{Z^2}{2} \left(\frac{\frac{(2)^2}{u^2}}{\frac{(1)}{u^4}} - \frac{1}{6} \frac{\frac{(3)}{u}}{\frac{(1)}{u^3}} - \frac{1}{12} \frac{B}{\frac{(1)}{u^2}} \right) + O(Z^3) \right]. \quad (7.9)$$

B. Classical cosmological tests

Classical cosmological tests, such as apparent magnitude – red shift ($m - Z$), number counts – red shift ($N - Z$), angular size – red shift relations, and some other, reveal specific

dependence of astrophysical observables on the angular coordinates (θ, ϕ) in a rotating world. Thus a careful analysis of the angular variations of empirical data over the whole celestial sphere is necessary.

The knowledge of null geodesics enables one to obtain the explicit form of the area distance r between an observer at a point P and any source S . Each ray (a null geodesic) $x^\mu(s; p^A)$ passing through P is labelled by the two parameters $p^A := (\theta, \phi)$, $A = 1, 2$ which have a simple meaning of the spherical angles fixing the direction of ray at P . Thus in general, r is a function of the direction of observation, that is $r = r(\theta, \phi)$. Besides this, r of course depends on the value of the affine parameter s , or equivalently on the moment of t at which source radiates a ray detected by an observer at t_0 . The area of the cross-section of the bundle of rays is

$$dA_S = \sqrt{\det G_{AB}} d\theta d\phi, \quad G_{AB} = g_{\mu\nu} \frac{\partial x^\mu}{\partial p^A} \frac{\partial x^\nu}{\partial p^B}. \quad (7.10)$$

Since the solid angle in (7.3) is defined, as usually, by $d\Omega_P = \sin \theta d\theta d\phi$, one easily obtains the general expression for the area distance between an observer P and a source S which lies on the geodesic labeled by the observational angles (θ, ϕ) at the value s of the geodetic parameter,

$$r(s; \theta, \phi) = \frac{\sqrt{\det G_{AB}(s; \theta, \phi)}}{\sin \theta}. \quad (7.11)$$

The knowledge of the null geodesics (6.5)-(6.8) makes it possible to calculate area distances in the Gödel type universe explicitly. For example, using (6.5)-(6.8), one can find for the area distance *along the axis of rotation* the following exact result

$$r^2(t; \theta = 0) = \frac{\sin^2 \left(\int_{t_0}^t dt' \omega(t') \right)}{\omega^2(t)}. \quad (7.12)$$

The rotational effects are maximal in the directions close to the z axis, and quite remarkably the observations in the direction of rotation can be described by simple and clear formulas.

As for an arbitrary direction, exact formulas become very complicated and it is much more convenient to replace them by the Kristian-Sachs expansions. One can use the expansions (7.8) and (7.9) in the calculations of observable effects in rotating cosmologies:

All the angular dependent rotational contributions are contained in the coefficients of these expansions.

For the Gödel type cosmology (4.1) the classical $(m - Z)$ and $(N - Z)$ relations read as follows [287,288,187].

Apparent magnitude m versus red shift Z : The apparent magnitude of a source S with the energy flux L_P , measured at P , is defined by $m = -\frac{5}{2} \log_{10} L_P$. Given the total output of the source L , $L_S = L/4\pi$, the luminosity distance is $D^2 := L_S/L_P = r^2(1 + Z)^4$, see [95,96]. Using (7.9), one finds for the Gödel type universe:

$$\begin{aligned} m = & M - 5 \log_{10} H_0 + 5 \log_{10} Z + \frac{5}{2} (\log_{10} e)(1 - q_0)Z \\ & - 5 \log_{10} \left(1 + \sqrt{\frac{\sigma}{k + \sigma}} \sin \theta \sin \phi \right) \\ & - \frac{5}{2} (\log_{10} e) \frac{\omega_0}{H_0} Z \sin \theta \cos \phi \left[\frac{\sqrt{\frac{\sigma}{k + \sigma}} + \sin \theta \sin \phi}{\left(1 + \sqrt{\frac{\sigma}{k + \sigma}} \sin \theta \sin \phi \right)^2} \right] + O(Z^2). \end{aligned} \quad (7.13)$$

Here $M = -\frac{5}{2} \log_{10} L_S$ is the absolute magnitude of a source with an intrinsic luminosity L_S .

Number of sources N versus red shift Z : Let us assume that the sources of electromagnetic radiation are distributed homogeneously with the average density $n(t)$. Neglecting the evolution of sources, one can count the number of images seen by an observer as follows. Consider the bundle with a solid angle $d\Omega$ detected by an observer at P . When moving along the central ray, the distance between the two infinitesimally close cross-sections is equal $d\ell = (k^\mu u_\mu) ds$, and the volume between these sections is $dA d\ell$, with $dA = r^2 d\Omega$. Hence, the number of sources contained between these two sections is given by $dN = -n_S (k^\mu u_\mu)_S r^2 d\Omega ds$. Integrating, and using (7.8)-(7.9):

$$\begin{aligned} \frac{dN}{d\Omega} = & \frac{n_0 Z^3}{3H_0^3 \left(1 + \sqrt{\frac{\sigma}{k + \sigma}} \sin \theta \sin \phi \right)^3} \left[1 - \frac{3}{2} (1 + q_0) Z \right. \\ & \left. - 3 \frac{\omega_0}{H_0} \frac{\sin \theta \cos \phi \left(\sqrt{\frac{\sigma}{k + \sigma}} + \sin \theta \sin \phi \right)}{\left(1 + \sqrt{\frac{\sigma}{k + \sigma}} \sin \theta \sin \phi \right)^2} Z + O(Z^2) \right]. \end{aligned} \quad (7.14)$$

The $(N - Z)$ relation (7.14) describes the number of sources per solid angle $d\Omega$ observed up to the value Z of red shift. One can estimate the global difference of the number of sources visible in two hemispheres of the sky, N^+, N^- , by integrating (7.14). The result is

$$\frac{N^+ - N^-}{N^+ + N^-} = \frac{1}{2} \sqrt{\frac{\sigma}{k + \sigma}} \left(3 - \frac{\sigma}{k + \sigma} \right) + O(Z^2). \quad (7.15)$$

It seems worthwhile to draw attention to the absence of a correction proportional to Z in (7.15). It is difficult to make a comparison of our results with [431,265,103] as they study stationary rotating models in which there is no red shift.

For some time classical cosmological tests were carefully carried out for standard models, but later it was recognized that evolution of physical properties of sources often dominates over geometrical effects. However, specific angular irregularities predicted in rotating cosmologies, (7.13), (7.14)-(7.15), may revive the importance of the classical tests.

C. Periodic structure of the universe

Recent analysis of the large-scale distribution of galaxies [54] has revealed an apparently periodic structure of the number of sources as a function of the red shift. Cosmic rotation may give a natural explanation of this fact [193]. The crucial point is the helicoidal behaviour of the null geodesics (6.5)-(6.7) in the Gödel type model in the directions close to the rotation axis. This yields a periodicity of the area distance as a function of red shift, and hence the visible distribution of sources turns out to be also approximately periodical in Z .

This effect is most transparent for the direction of rays straight along the axis of rotation. The area distance is then given by (7.12). In order to be able to make some quantitative estimates, let us assume the polynomial law for the scale factor,

$$a(t) = a_0 \left(\frac{t - t_\infty}{t_0 - t_\infty} \right)^\gamma, \quad (7.16)$$

which is naturally arising in a number of cosmological scenarios ($0 < \gamma < 1$). Then one can derive (analogously to (7.14)) the distribution of number of sources per red shift per solid angle:

$$\frac{dN}{d\Omega dZ} = \frac{n_0}{\omega_0^2 H_0 (1+Z)^{1/\gamma}} \sin^2 \left(\frac{\gamma \omega_0}{(1-\gamma) H_0} \left[(1+Z)^{(\gamma-1)/\gamma} - 1 \right] \right). \quad (7.17)$$

Thus, the apparent distribution of visible sources is an oscillating function of red shift, with slowly decreasing amplitude. A similar generalized formula can be obtained for arbitrary directions, so that (7.17) is modified by additional angular dependence of the magnitude of successive extrema of distribution function.

The observational data [54] give for the distance between maxima the value $128h^{-1}$ Mps (where $H_0 = 100h \text{ km sec}^{-1} \text{ Mps}^{-1}$). From this one can estimate the rotation velocity which is necessary to produce such a periodicity effect,

$$\omega_0 \approx 74H_0. \quad (7.18)$$

This result does not depend on γ .

D. Polarization rotation effect

The cosmic rotation affects a polarization of radiation which propagates in the curved spacetime (4.1), and this produces a new observable effect which has been reported in the literature by Birch [46,47] and more recently by Nodland and Ralston [279,280]. In the geometrical optics approximation, polarization is described by a space-like vector f^μ which is orthogonal to the wave vector, $f_\mu k^\mu = 0$, and is parallelly transported along the light ray,

$$k^\mu \nabla_\mu f^\nu = 0. \quad (7.19)$$

The influence of the gravitational field of a compact rotating massive body on the polarization of light was investigated by many authors, see e.g., [399,426,26,340,18,20,53,101,115,179,258]. The analysis of (7.19) for the rotating cosmologies shows that the global cosmic rotation also forces a polarization vector to change its orientation during the propagation along a null geodesics [287,257,216]. It is clear that this conclusion has physical meaning only when one can define a frame at any point of the ray with respect to which polarization rotates. Let us describe how this can be achieved.

As it is well known, gravitational field affects all the properties of an image of a source, such as shape, size and orientation [377,286,95,96,389]. Like the rotation of the polarization vector, the deformation and rotation of an image depend both on the local coordinates and on the choice of an observer's frame of reference. However, if one considers a combination of the two problems, then a truly observable effect arises which is coordinate and frame independent. Putting it in another way, one should calculate the influence of the cosmic rotation on *the relative angle* η between the polarization vector and the direction of a major axis of an image [194]. This problem was discussed in the papers [314,141], but in our opinion, their results are incomplete, in the sense that only null congruences converging at the point of observation were considered.

The relative angle η can be most conveniently defined within the framework of the Newman-Penrose spin coefficient formalism. Namely, it is sufficient to construct a null frame $\{l, n, m, \bar{m}\}$ is such a way that l coincides with the wave vector k , and the rest of the vectors are covariantly constant along l . Then we can consider m as a polarization vector, and thus a deformation of an image of a source, calculated with respect to the frame $\{l, n, m, \bar{m}\}$, gives the observable relative angle η . Let us describe explicitly the null frame:

$$l = \frac{a_0}{a} \left[\left\{ 1 + \sqrt{\frac{\sigma}{k + \sigma}} \sin \theta \sin \Phi \right\} \partial_t + \frac{1}{a} \sin \theta \cos \Phi \partial_x + \frac{e^{-mx}}{a\sqrt{k + \sigma}} \sin \theta \sin \Phi \partial_y + \frac{1}{a} \cos \theta \partial_z \right], \quad (7.20)$$

$$n = \frac{a}{a_0} \left(\frac{1}{1 + \cos \theta} \right) \left[\partial_t - \frac{1}{a} \partial_z \right], \quad (7.21)$$

$$m = \frac{e^{i\Psi}}{\sqrt{2}} \left[\left\{ \sqrt{\frac{\sigma}{k + \sigma}} + i \left(\frac{\sin \theta}{1 + \cos \theta} \right) e^{-i\Phi} \right\} \partial_t + \frac{i}{a} \partial_x + \frac{e^{-mx}}{a\sqrt{k + \sigma}} \partial_y - \frac{i}{a} \left(\frac{\sin \theta}{1 + \cos \theta} \right) e^{-i\Phi} \partial_z \right], \quad (7.22)$$

$$\bar{m} = \frac{e^{-i\Psi}}{\sqrt{2}} \left[\left\{ \sqrt{\frac{\sigma}{k + \sigma}} - i \left(\frac{\sin \theta}{1 + \cos \theta} \right) e^{i\Phi} \right\} \partial_t - \frac{i}{a} \partial_x + \frac{e^{-mx}}{a\sqrt{k + \sigma}} \partial_y + \frac{i}{a} \left(\frac{\sin \theta}{1 + \cos \theta} \right) e^{i\Phi} \partial_z \right]. \quad (7.23)$$

Here

$$\Psi(t, z) = z \frac{m}{2} \sqrt{\frac{\sigma}{k + \sigma}} + \Phi(t). \quad (7.24)$$

One should note that (7.20)-(7.23) is a smooth field of frames which covers the whole space-time manifold. Direct computation proves that (7.20) is the null geodesic congruence with an affine parametrization. We say that this congruence is oriented along a direction given by the spherical angles (θ, ϕ) in the local Lorentz frame of an observer at P , because a null geodesics (6.5)-(6.8) belongs to this congruence.

It is straightforward to find the spin coefficients (we are using the definitions of [71], and mark the spin coefficients with tildes in order to distinguish them from the other quantities in our paper):

$$\tilde{\varepsilon} = 0, \quad \tilde{\kappa} = 0, \quad \tilde{\lambda} = 0, \quad \tilde{\nu} = 0, \quad (7.25)$$

$$\begin{aligned} \tilde{\rho} = & -\frac{a_0}{a^2} \left[\dot{a} \left\{ 1 + \sqrt{\frac{\sigma}{k + \sigma}} \sin \theta \sin \Phi \right\} + \frac{m}{2} \left(\frac{k}{k + \sigma} \right) \frac{\sin \theta \cos \Phi}{(1 + \sqrt{\frac{\sigma}{k + \sigma}} \sin \theta \sin \Phi)} \right] \\ & + i \frac{a_0}{a^2} \frac{m}{2} \cos \theta \left(\frac{\sqrt{\frac{\sigma}{k + \sigma}} + \sin \theta \sin \Phi}{1 + \sqrt{\frac{\sigma}{k + \sigma}} \sin \theta \sin \Phi} \right), \end{aligned} \quad (7.26)$$

$$\begin{aligned} \tilde{\sigma} = & \frac{a_0}{a^2} \frac{m}{2} \left(\frac{k}{k + \sigma} \right) \frac{e^{2i\Psi} \sin \theta}{(1 + \sqrt{\frac{\sigma}{k + \sigma}} \sin \theta \sin \Phi)} \left(-\cos \Phi [2 \cos \theta - 1 \right. \\ & \left. - 2 \cos^2 \Phi (\cos \theta - 1)] + i \sin \Phi [\cos \theta - 2 \cos^2 \Phi (\cos \theta - 1)] \right). \end{aligned} \quad (7.27)$$

We do not write the other spin coefficients, because their values are irrelevant. Only one important step remains to be done: the spin coefficient $\tilde{\pi} \neq 0$ in the frame (7.20)-(7.23), and we need to make an additional Lorentz transformation

$$l \longrightarrow l, \quad n \longrightarrow n + A^* m + A \bar{m} + A^* A l, \quad m \longrightarrow m + A l, \quad \bar{m} \longrightarrow \bar{m} + A^* l, \quad (7.28)$$

where the function A satisfies equation $l^\mu \partial_\mu A + \tilde{\pi}^* = 0$. This ensures that $\tilde{\pi} = 0$ in a new frame, while remarkably the transformation (7.28) does not change any of the spin coefficients (7.25)-(7.27). Thus one obtains finally the field of null frames $\{l, n, m, \bar{m}\}$ with the required properties: l is the null geodesic congruence with affine parametrization, while $n, m, \bar{m}$ are covariantly constant along l . The latter is equivalent to $\tilde{\kappa} = \tilde{\varepsilon} = \tilde{\pi} = 0$.

As it is well known, deformation and rotation of an image along a null geodesics are described by the optical scalars [377]:

$$\tilde{\vartheta} = -\operatorname{Re}\tilde{\rho}, \quad \tilde{\omega} = \operatorname{Im}\tilde{\rho}, \quad \tilde{\sigma}. \quad (7.29)$$

We now assume, for definiteness, that the polarization vector f^μ coincides with the vector m^μ of the above null frame. Let an image of a source, as seen at the point corresponding to a value $s = s_1$ of the affine parameter, be an ellipse with the major axis p and the minor axis q . Then one can straightforwardly obtain [194] for the angle of rotation of the major axis of the image at $s_2 = s_1 + \delta s$,

$$\delta\eta = -\tilde{\omega}\delta s - \frac{p^2 + q^2}{p^2 - q^2} \operatorname{Im}\tilde{\sigma} \delta s. \quad (7.30)$$

Integration along a ray gives the finite angle of rotation.

It is worthwhile to notice that exactly along the cosmic rotation axis the observer at P finds for the optical scalars

$$\tilde{\vartheta}_P = H_0, \quad \tilde{\omega}_P = \omega_0, \quad \tilde{\sigma} = 0. \quad (7.31)$$

Thus the effect of rotation of the polarization vector in this direction is the most explicit.

As for an arbitrary direction of observation, we finally find from (7.30) for $k/\sigma \ll 1$ with the help of the Kristian–Sachs expansions (7.7)–(7.9):

$$\eta = \omega_0 r \cos\theta + O(Z^2). \quad (7.32)$$

This result is in a good agreement with the observational data reported [46] on the dipole anisotropy of distribution of the difference between the position angles of elongation (the major axis) and polarization in a sample of 3CR radio sources. The estimate for the direction and the magnitude of the cosmic rotation, obtained in [288,187,194] from Birch’s data, reads

$$l^\circ = 295^\circ \pm 25^\circ, \quad b^\circ = 24^\circ \pm 20^\circ, \quad (7.33)$$

$$\omega_0 = (1.8 \pm 0.8)H_0. \quad (7.34)$$

Not entering into the discussion [428,94,229,68] of the statistical significance of the results of Nodland and Ralston, we can make a comparison with the new data. In [279], the dipole effect of the rotation of the plane of polarization was reported in the form $\beta = \frac{1}{2}\Lambda_s^{-1} r \cos\gamma$, where β is the residual rotation angle between the polarization vector and the direction of the major axis of a source (remaining after the Faraday rotation is extracted), r is the distance to the source, γ is the angle between the direction of the wave propagation and the constant vector $\vec{s}$. The analysis of the data for 160 radio sources yielded [279] the best fit for the constant $\Lambda_s = (1.1 \pm 0.08) \frac{2}{3} \frac{10^{15}}{H} \text{ m yr}^{-1}$ (with the Hubble constant H), and the $\vec{s}$ -direction RA (21h $\pm$ 2h), dec ($0^\circ \pm 20^\circ$). In an attempt to explain their observations, Nodland and Ralston concluded that it is impossible to understand such an effect within the conventional physics. Instead, they considered a modified electrodynamics with the Chern-Simons type term violating the Lorentz invariance [67], and related Λ_s to the coupling constant of that term. However, interpreting the effect as arising from the cosmic rotation, we immediately find

$$l^\circ = 50^\circ \pm 20^\circ, \quad b^\circ = -30^\circ \pm 25^\circ, \quad (7.35)$$

$$\omega_0 = (6.5 \pm 0.5) H_0. \quad (7.36)$$

This is larger than the estimate (7.34) obtained from Birch's data [46]. Also the direction of the cosmic vorticity is different from (7.33). However, it is interesting to note that within the error limits, the two directions are orthogonal to each other.

It is clear that further careful observations and statistical analyzes will be extremely important in establishing the true value of the cosmic rotation. In particular, it has been recently claimed in [68], that the magnitude of the polarization rotation effect should be one or two orders lower than reported in [279]. The value of the cosmic vorticity then should be reduced correspondingly.

8. CONCLUSIONS

In our discussion of the properties of rotating cosmologies we have paid special attention to the Gödel type model (4.1). However the main conclusions are true also for the whole class of metrics (3.1). One can notice, that in some cases the numerical estimates obtained for the value of the vorticity do not agree with each other, e.g., see (7.18), (7.34), and (7.36). Such inconsistencies can reflect the fact that too few empirical data were analyzed until now, so that a further detailed discussion is required in order to obtain the final convincing estimates. Already now certain modifications of the above simple models are necessary: It is clear, in the light of the modern COBE results [402,41,436], that the purely rotating models (3.1) should be replaced by the cosmologies with a nontrivial shear. A preliminary analysis of such generalizations shows that rotating models can be made compatible with the COBE data without destroying the rest of the rotational effects (in particular, without essential modification of the polarization rotation formulas). It may turn out, though, that some (or all) of the above mentioned effects are explained after all by some different physical (and geometrical) reasons, not related to the cosmic rotation. However, “whether our universe is rotating or not, it is of fundamental interest to understand the interrelation between rotation and other aspects of cosmological models as well as to understand the observational significance of an overall rotation” [369].

We believe that the cosmic rotation is an important physical effect which should find its final place in cosmology. In this paper we outlined one of the possible theoretical frameworks which can underlie our understanding of this phenomenon.

Acknowledgments. This work was partly supported by the Deutsche Forschungsgemeinschaft (Bonn) grant He 528/17-2. I thank Prof. K.-E. Hellwig for the kind invitation to the Colloquium on Cosmic Rotation at the Technische Universität Berlin. The warm hospitality of the organizers and highly interesting discussions with the participants are gratefully acknowledged.

REFERENCES

- [1] A.J. Accioly, *An unusual cosmological solution in the context of higher-derivative gravity*, *Nuovo Cim.* **B100** (1987) 703-707.
- [2] A.J. Accioly, *Is higher-derivative gravity a good therapy for the causal pathologies of Gödel-type universes?*, *Progr. Theor. Phys.* **79** (1988) 1269-1272.
- [3] A.J. Accioly and A.T. Gonçalves, *Homogeneous space-times of Gödel-type in higher-derivative gravity*, *J. Math. Phys.* **28** (1987) 1547-1552.
- [4] A.J. Accioly and A.T. Gonçalves, *Can the causal pathologies of Gödel-type universes be avoided in higher-derivative gravity?*, Preprint IFUSP/P-605 (Univ. Sao Paulo, October 1986) 17 p.
- [5] A.J. Accioly and G.E.A. Matsas, *Are there causal vacuum solutions with symmetries of the Gödel universe in higher-derivative gravity?*, *Phys. Rev.* **D38** (1988) 1081-1084.
- [6] R. Adler, M. Bazin, and M. Schiffer, *Introduction to general relativity* (McGraw-Hill: New York, 1975).
- [7] V.G. Agakov, *Physically homogeneous spaces with rotation*, *Vestnik Mosk. Univ., ser. fizika, astronomiya*, **20**, n. 4 (1979) 73-75 (in Russian).
- [8] V.G. Agakov, *The nonstationary generalization of the Gödel cosmological model*, *Gen. Rel. Grav.* **16** (1984) 317-323.
- [9] V.G. Agakov, *Homogeneous cosmological models with rotation*, in: *Abstracts of Contr. Papers at the 7th Gravit. Soviet Conf. (Erevan, 1988)* (Erevan State Univ. Press: Erevan, 1988) 397-398 (in Russian).
- [10] D.V. Ahluwalia and T. Goldman, *Space-time symmetries and vortices in the cosmos*, *Mod. Phys. Lett.* **A8** (1993) 2623-2630.
- [11] E.S. Akeley, *The axially symmetric stationary gravitational field*, *Phil. Mag.* **11** (1931)

322-330.

- [12] E.S. Akeley, *The rotating fluid in the relativity theory*, *Phil. Mag.* **11** (1931) 330-344.
- [13] R.A. Alper, H. Bete, and G. Gamow, *The origin of chemical elements*, *Phys. Rev.* **73** (1948) 803-804.
- [14] H. Arp, *The 37.5 km s^{-1} redshift periodicity of galaxies as the machion frequency*, *Phys. Lett.* **B387** (1996) 48-50.
- [15] J.E. Åman, J.B. Fonseca-Neto, M.A.H. MacCallum and M.J. Rebouças, *Riemann-Cartan space-times of Gödel type*, *Class. Quantum Grav.* **15** (1998) 1089-1101.
- [16] R.R. Andreev, *The problem of the existence of large-scale anisotropy in metagalactic space*, *Astrophysics (USA)* **24** (1986) 213-220 [*Astrofizika* **24** (1986) 363-375 (in Russian)].
- [17] R.R. Andreev, T.G. Arshakyan, A.N. Makarov, and M.A. Mnatsakanyan, *On the large-scale anisotropy of metagalaxy*, in: *Abstracts of Contr. Papers at the 7th Gravit. Soviet Conf. (Erevan,1988)* (Erevan State Univ. Press: Erevan, 1988) 398-400 (in Russian).
- [18] A.M. Anile and R.A. Breuer, *Gravitational Stokes parameters*, *Astrophys. J.* **189** (1974) 39-49.
- [19] A.M. Anile and S. Motta, *Vorticity perturbations and isotropy of the cosmic microwave background*, *Astron. and Astrophys.* **49** (1976) 205-209.
- [20] A.M. Anile and S. Motta, *Perturbations of the general Robertson-Walker universes and angular variations of the cosmic blackbody radiation*, *Astrophys. J.* **207** (1976) 685-692.
- [21] A.M. Anile and R.A. Breuer, *Polarization transport in anisotropic universes*, *Astrophys. J.* **217** (1977) 353-357.
- [22] A.M. Anile, L. Danese, G. De Zotti, and S. Motta, *Cosmological turbulence reexamined*,

- Astrophys. J.* **205** (1976) L59-L63.
- [23] H. Asada and M. Kasai, *Lagrangian description of fluid flow with vorticity in relativistic cosmology*, *Phys. Rev.* **D59** (1999) 123515(5).
- [24] E. Asseo and H. Sol, *Extragalactic magnetic fields*, *Phys. Rept.* **148** (1987) 307-436.
- [25] S. Bajtlik, R. Juszkiewicz, M. Prószyński, and P. Amsterdamsky, *2.7K radiation and the isotropy of the universe*, *Astrophys. J.* **300** (1986) 463-473.
- [26] N.L. Balazs, *Effect of a gravitational field, due to a rotating body, on the plane of polarization of electromagnetic wave*, *Phys. Rev.* **110** (1958) 236-239.
- [27] F. Bampi and C. Zordan, *A note on Gödel's metric*, *Gen. Rel. Grav.* **9** (1978) 393-398.
- [28] A. Banerjee and S. Banerji, *Stationary distributions of dust and electromagnetic fields in general relativity*, *J. Phys.* **A1** (1968) 188-193.
- [29] J.D. Barrow, *General relativistic cosmological models and the cosmic microwave background radiation*, *Canad. J. Phys.* **64** (1986) 152-159.
- [30] J.D. Barrow and M.P. Dabrowski, *Gödel universes in string theory*, *Phys. Rev.* **D58** (1998) 103502 (5 pages).
- [31] J.D. Barrow and K.E. Kunze, *String cosmology*, *Preprint Univ. Sussex* (July, 1998) 12 p., e-Archive gr-qc/9807040.
- [32] J.D. Barrow, R. Juszkiewicz, and D.H. Sonoda, *Structure of the cosmic microwave background*, *Nature* **305** (1983) 397-402.
- [33] J.D. Barrow, R. Juszkiewicz, and D.H. Sonoda, *Universal rotation: how large can it be?*, *Mon. Not. R. Astr. Soc.* **213** (1985) 917-943.
- [34] N. Batakis, *Imparting small vorticity to a Bianchi type- VI_h empty spacetime*, *Phys. Rev.* **D23** (1981) 1681-1682.

- [35] N. Batakis, *Electromagnetic field generated by cosmic vorticity*, *Phys. Lett.* **A85** (1981) 409-410.
- [36] N. Batakis and J.M. Cohen, *Cosmological model with expansion, shear and vorticity*, *Phys. Rev.* **D12** (1975) 1544-1550.
- [37] S.S. Bayin and F.I. Cooperstock, *Rotational perturbations of Friedman universes*, *Phys. Rev.* **D22** (1980) 2317-2322.
- [38] S.S. Bayin, *Slowly rotating fluid spheres in general relativity with and without radiation*, *Phys. Rev.* **D24** (1981) 2056-2065.
- [39] S.S. Bayin, *Slowly rotating fluid spheres in general relativity*, *Phys. Rev.* **D27** (1983) 3030-3031.
- [40] M. Bedran, V. Vaidya, and M. Som, *Stationary cosmological solutions with torsion*, *Nuovo Cim.* **B87** (1985) 101-108.
- [41] C.L. Bennett et al, *Preliminary separation of galactic and cosmic microwave emission for the COBE Differential Microwave Radiometer*, *Astrophys. J.* **396** (1992) L7-12.
- [42] R. Bergamini and G. Giampieri, *Influence of rotation on the initial state of the universe*, *Phys. Rev.* **D40** 3960-3967.
- [43] M.F. Bietenholz and P.P. Kronberg, *Is there really evidence for the universal rotation?*, *Astrophys. J.* **287** (1984) L1-2.
- [44] M.F. Bietenholz, *Determining dependencies between directional quantities and position on a sphere*, *Astron. J.* **91** (1986) 1249-1252.
- [45] S. Bildhauer, *Freely propagating polarized radiation in curved space-times*, *Astron. Astrophys.* **219** (1989) 25-28.
- [46] P. Birch, *Is the universe rotating?*, *Nature* **298** (1982) 451-454.

- [47] P. Birch, *Is there evidence for universal rotation? Birch replies*, *Nature* **301** (1982) 736.
- [48] G. Boerner, *The early universe - facts and fiction*, *Texts and Monographs In Physics* (Springer: Berlin, 1993).
- [49] A. Braccesi, *Inflation and Mach's principle*, *Astron. Astrophys.* **194** (1988) 1-2.
- [50] R. N. Bracewell and V. R. Eshleman, *The Nodland-Ralston effect, or the up and down of the universe*, *Preprint of Space, Telecommunications and Radio-science Laboratory, Stanford, CA 94305* (April 24, 1997); available at <http://www-star.stanford.edu/~bracewel/nreff.html>.
- [51] J. Bradley and E. Sviestins, *Some rotating, time-dependent Bianchi type VIII cosmologies with heat flow*, *Gen. Rel. Grav.* **16** (1984) 1119-1133.
- [52] M. Bradley, G. Fodor, L.A. Gergely, M. Marklund, and Z. Perjés, *Rotating perfect fluid sources of the NUT metric*, *Preprint KFKI Budapest* (1998) 8 p.; e-print Archive: gr-qc/9807058.
- [53] C. Brans, *Propagation of electromagnetic polarization effects in anisotropic cosmologies*, *Astrophys. J.* **197** (1975) 1-4.
- [54] T.J. Broadhurst, R.S. Ellis, D.G. Koo, and A.S. Szalay, *Large-scale distribution of galaxies at the Galactic poles*, *Nature* **343** (1990) 726-728.
- [55] T.J. Broadhurst, R.S. Ellis, and T. Shanks, *The Durham/Anglo-Australian telescope faint galaxy redshift survey*, *Mon. Not. R. Astr. Soc.* **235** (1988) 827-856.
- [56] K.A. Bronnikov, Yu.P. Rybakov, and G.N. Shikin, *Cylindrical solitons in a Gödel universe and their stability*, *Sov. Phys. J.* **34** n. 5 (1991) 404-406.
- [57] P. Brosche, *The mass – angular momentum – diagram of astronomical objects*, in: “*Proc. of the 6th Course of Internat. School on Cosmology and Gravitation: Spin,*

- Torsion, Rotation and Supergravity*” (Erice, 1979) Eds. P.G.Bergmann and V.De Sabata (Plenum: New York, 1980) 375-382.
- [58] P. Brosche, *Angular momentum versus mass, Comments on Astrophysics* **11** (1986) 213-221.
- [59] P. Brosche and L.J. Tassie, *A scenario for the formation of astronomical objects from superstrings, Astron. Astrophys.* **219** (1989) 13-24.
- [60] P. Brosche and L.J. Tassie, *Hierarchical dark matter from fragmenting macroscopic superstrings, Astron. Nachr.* **316** (1995) 149-153.
- [61] E.F. Bunn, P.G. Ferreira, and J. Silk, *How anisotropic is our universe, Phys. Rev. Lett.* **77** (1996) 2883-2886.
- [62] M.O. Calvão, M.J. Rebouças, A.F.F. Teixeira, and W.M. Silva, Jr., *Notes on a class of homogeneous space-times, J. Math. Phys.* **29** (1988) 1127-1131.
- [63] S. Carneiro, *The large numbers hypothesis and quantum mechanics, Found. Phys. Lett.* **11** (1998) 95-102.
- [64] S. Carneiro, *A Gödel-Friedmann cosmology?, Phys. Rev.* **D61** (2000) 083506 (5 pages).
- [65] S. Carneiro, *Open cosmologies with rotation, Preprint IMFF, Madrid (June, 2000) 15 p.* E-print Archive: gr-qc/0003096.
- [66] H.L. Carrion, M.J. Rebouças, and A.F.F. Teixeira, *Gödel-type space-times in induced matter gravity theory, J. Math. Phys.* **40** (1999) 4011-4027.
- [67] S.M. Carroll, G.B. Field, and R. Jackiw, *Limits on a Lorentz- and parity-violating modifications of electrodynamics, Phys. Rev.* **D41** (1990) 1231-1240.
- [68] S.M. Carroll and G.B. Field, *Is there evidence for cosmic anisotropy in the polarization of distant radio sources?, Phys. Rev. Lett.* **79** (1997) 2394-2397.

- [69] S.K. Chakraborty, *A remark on a cylindrically symmetric rotating metric*, *Gen. Rel. Grav.* **12** (1980) 925-928.
- [70] S.K. Chakraborty and N. Bandyopadhyay, *Gödel-type universe with perfect fluid and a scalar field*, *J. Math. Phys.* **24** (1983) 129-132.
- [71] S. Chandrasekhar, *The mathematical theory of black holes* (Clarendon Press: Oxford, 1983).
- [72] S. Chandrasekhar and J.P. Wright, *The geodesics in Gödel's universe*, *Proc. Nat. Acad. Sci. USA* **47** (1961) 341-347.
- [73] A.D. Chernin, *Origin of rotation of galaxies*, *Astrofizika* **10** (1974) 535-542 (in Russian).
- [74] A.D. Chernin, *Origin of rotation of galaxies. II*, *Astrofizika* **13** (1977) 69-78 (in Russian).
- [75] A.D. Chernin, *On the origin of proto-galactic eddies*, *Astrophys. Lett.* **19** (1978) 97-100.
- [76] A.D. Chernin, *Eddies in the Universe*, *Riv. Nuovo Cim.* **19** (1996) 1-98.
- [77] A.D. Chernin and A.Yu. Ushakov, *Angular momentum of galaxies and the structure of protogalactic vorticity*, *Astrophys. Space Sci.* **96** (1983) 217-221.
- [78] L.P. Chimento, A.S. Jakubi, and J. Pullin, *Coleman-Weinberg symmetry breaking in a rotating spacetime*, *Class. Quantum Grav.* **6** (1989) L45-L48.
- [79] J.M. Cohen, C.V. Vishveshwara, and S.V. Dhurandbar, *Electromagnetic fields in the Gödel universe*, *J. Phys.: Math. and Gen.* **A13** (1980) 933-938.
- [80] P. Coles, *Understanding recent observations of the large-scale structure of the universe*, *Nature* **346** (1990) 446-447.

- [81] C.B. Collins and S.W. Hawking, *The rotation and distortion of the universe*, *Mon. Not. R. Astr. Soc.* **162** (1973) 307-320.
- [82] R.G. Conway, P. Birch, R.J. Davis, L.R. Jones, A.J. Kerr, and D. Stannard, *Multi-frequency observations of linear polarization in 94 sources from the 3CR catalogue. I. Observations*, *Mon. Not. R. Astr. Soc.* **202** (1983) 813-823.
- [83] N. Dallaporta and F. Luccin, *On galaxy formation from primeval universal turbulence*, *Astron. and Astrophys.* **19** (1972) 123-134.
- [84] N. Dallaporta and F. Luccin, *On galaxy formation from primeval universal turbulence. II*, *Astron. and Astrophys.* **26** (1973) 325-333.
- [85] I. Damião Soares, *Inhomogeneous rotating universes with closed timelike geodesics of matter*, *J. Math. Phys.* **21** (1980) 521-525.
- [86] P. Davies, *Does the universe rotate?*, *Sky & Telesc.* **75** (1988) 599-602.
- [87] H. Dehnen and H. Hönl, *Finite universe and Mach's principle*, *Nature* **106**, n. 4852 (1962) 362-363.
- [88] M. Demiański and L.P. Grishchuk, *Homogeneous rotating universe with flat space*, *Commun. Math. Phys.* **25** (1972) 233-244.
- [89] A. Dobado and A.L. Maroto, *Primordial torsion fields as an explanation of the anisotropy in cosmological electromagnetic propagation*, *Mod. Phys. Lett.* **A12** (1997) 3003-3007.
- [90] J. Duarte de Oliveira, A.F.F. Teixeira, and J. Tiomno, *Homogeneous cosmos of Weyssenhoff fluid in Einstein-Cartan space*, *Phys. Rev.* **D34** (1986) 3661-3665.
- [91] K. Dunn, *Two-fluid cosmological models in Gödel-type spacetimes*, *Gen. Rel. Grav.* **21** (1989) 137-147.
- [92] J. Ehlers, *Contributions to the relativistic mechanics of continuous media*, *Gen. Rel.*

- Grav.* **25** (1993) 1225-1266 [English version of: *Beiträge zur relativistischen Mechanik kontinuierlicher Medien*, Akad. der Wiss. und der Lit. Mainz, Abh. der math. naturwiss. Kl. n. **11** (1961) 791-837].
- [93] J. Einasto, M. Einasto, S. Gottlober, V. Muller, V. Saar, A.A. Starobinsky, E. Tago, D. Tucker, H. Andernach, and P. Frisch, *A 120-Mpc periodicity in the three-dimensional distribution of galaxy superclusters*, *Nature* **385** (1997) 139-141.
- [94] D.J. Eisenstein and E.F. Bunn, *Comment on the appropriate null hypothesis for cosmological birefringence*, *Phys. Rev. Lett.* **79** (1997) 1957-1958.
- [95] G.F.R. Ellis, *Relativistic cosmology*, in: *General Relativity and Cosmology* (Rendiconti S.I.F., Corso XLVIII, Varenna, 1969), Ed. R.K. Sachs (Academic Press: New York, 1971) 104-182.
- [96] G.F.R. Ellis, *Relativistic cosmology*, in: *Cargèse Lect. in Phys.* vol. **6**, Ed. E. Schatzman (Gordon and Breach: NY, 1973) 1-60.
- [97] G.F.R. Ellis, *The expanding Universe: A history of cosmology from 1917 to 1960*, in: *"Einstein and the history of general relativity"*, Eds. D. Howard and J. Stachel (Birkhäuser: Boston, 1986) 367-431.
- [98] J. Ellis and K. Olive, *Inflation can solve the rotation problem*, *Nature* **303** (1983) 679-681.
- [99] D.L. Farnsworth and R.P. Kerr, *Homogeneous dust-filled cosmological solutions*, *J. Math. Phys.* **7** (1966) 1625.
- [100] I. Farup and Ø. Grøn, *Vacuum energy and inertial dragging*, *Gen. Rel. Grav.* **28** (1996) 441-449.
- [101] F. Fayos and J. Llosa, *Gravitational effects on the polarization plane*, *Gen. Rel. Grav.* **14** (1982) 965-978.

- [102] A.J. Fennelly, *Gravitational charge, hadron hydrodynamics and Gödelised hadrons*, *Nature* **248** (1974) 221-223.
- [103] A.J. Fennelly, *Effects of a rotation of the universe on the number counts of radio sources: Gödel's universe*, *Astrophys. J.* **207** (1976) 693-699.
- [104] A.J. Fennelly, *Expanding shearfree spatially homogeneous universes with a nonsynchronous time coordinate and anisotropy of the universe*, *J. Math. Phys.* **19** (1978) 158-163.
- [105] A.J. Fennelly, *Observer frame rotation rates and magnetic fields in spatially homogeneous universes*, *J. Math. Phys.* **22** (1981) 126-132.
- [106] A.J. Fennelly, *Some effects of magnetic fields in spatially homogeneous universes with conductivity*, *Phys. Rev.* **D21** (1980) 2107-2118.
- [107] A.J. Fennelly, *Observer reference triad rotation, magnetic fields, and rotation in Euclidean cosmological models*, *Astrophys. J.* **252** (1982) 410-417.
- [108] A.J. Fennelly and C.R. Evans, *Magnetohydrodynamic perturbations of Robertson-Walker universes and of anisotropic Bianchi type-I universes*, *Nuovo Cim.* **B60** (1980) 1-45.
- [109] B.D.B. Figueredo, *Gödel-type spacetimes and the motions of charged particles*, *Class. Quantum Grav.* **15** (1998) 3849-3862.
- [110] B.D.B. Figueredo, I. Damião Soares, and J. Tiomno, *Gravitational coupling of Klein-Gordon and Dirac particles to matter vorticity and spacetime torsion*, *Class. Quantum Grav.* **9** (1992) 1593-1617.
- [111] J.B. Fonseca-Neto and M.J. Rebouças, *On a class of Riemann-Cartan space-times of Gödel type*, *Gen. Rel. Grav.* **30** (1998) 1301-1317.
- [112] G. Gamow, *Rotating universe?*, *Nature* **158** (1946) 549.

- [113] G. Gamow, *The role of turbulence in the evolution of the universe*, *Phys. Rev.* **86** (1952) 251.
- [114] E.N. Glass, *The Weyl tensor and shear-free perfect fluids*, *J. Math. Phys.* **16** (1975) 2361-2363.
- [115] N.Yu. Gnedin and I.G. Dymnikova, *Rotation of the plane of polarization of a photon in a Petrov type D space-time*, *Sov. Phys. JETP* **67** (1988) 13-15 [*ZhETF* **94** no. 1 (1988) 26-31 (in Russian)].
- [116] K. Gödel, *An example of a new type of cosmological solutions of Einstein's field equations of gravitation*, *Rev. Mod. Phys.* **21** (1949) 447-450.
- [117] K. Gödel, *Rotating universes in general relativity theory*, in: *Proc. of Int. Congress of Mathematicians (Cambridge, USA, 1950)*, Eds. L.M. Graves, E. Hille, P.A. Smith, and O. Zariski, (American Math. Soc.: Providence, RI, 1952) v. **1**, 175-181.
- [118] K. Gödel, *A remark about the relationship between relativity theory and idealistic philosophy*, in: "Albert Einstein, philosopher-scientist", *Library of living philosophers*, vol. 7, Ed. P.A. Schlipp (Evanstone: Library of living philosophers, 1949) 555-562.
- [119] H. Goenner, *Einführung in die Kosmologie* (Spektrum Akad. Verlag: Heidelberg, 1994).
- [120] M. Goldhaber and V. Trimble, *Limits on the chirality of interstellar and intergalactic space*, *J. Astrophysics and Astronomy* **17** (1996) 17-21.
- [121] F.M. Gomide, M.S. Berman, and R.L. Garcia, *A cosmological model with cylindrical symmetry*, *Rev. Mex. Astron. Astrofis.* **12** (1986) 46-48.
- [122] J. Gribbin, *Quantized quasars and a spinning universe*, *New Sci.* **100** n. 1384 (1983) 526.
- [123] L.P. Grishchuk, *Cosmological rotation of the quantum-mechanical origin and anisotropy of the microwave background*, *Phys. Rev.* **D48** (1993) 5581-5593.

- [124] L.P. Grishchuk, A.G. Doroshkevich, and V.N. Lukash, *The model of “mixmaster universe” with arbitrarily moving matter*, *Sov. Phys. JETP*. **28** (1969) 1210-1215 [*ZhETF* **55** (1969) 2281-2290 (in Russian)].
- [125] L.P. Grishchuk, A.G. Doroshkevich, and I.D. Novikov, *Anisotropy of the early stages of cosmological expansion and of relict radiation*, *Sov. Phys. JETP*. **34** (1971) 1-7 [*ZhETF* **61** (1971) 3-14 (in Russian)].
- [126] Ø. Grøn, *Expansion isotropization during the inflationary era*, *Phys. Rev.* **D32** (1985) 2522-2527.
- [127] Ø. Grøn, *Transition of a rotating Bianchi type-IX cosmological model into an inflationary era*, *Phys. Rev.* **D33** (1986) 1204-1205.
- [128] Ø. Grøn and H.H. Soleng, *Decay of primordial cosmic rotation in inflationary cosmologies*, *Nature* **328** (1987) 501-503.
- [129] Ø. Grøn and H.H. Soleng, *Cosmic inertial dragging*, *Class. Quantum Grav.* **5** (1988) L101-L104.
- [130] Ø. Grøn and H.H. Soleng, *The Vaidya-Patel solution with Robertson-Walker metric as a rotating inflationary scenario*, *J. Math. Phys.* **29** (1988) 1514-1517; *Erratum*, *J. Math. Phys.* **29** (1988) 2320.
- [131] Ø. Grøn and H.H. Soleng, *Note on rotating universe models*, *Acta Phys. Pol.* **B20** (1989) 557.
- [132] S.W. Goode, *Vorticity and isotropic singularities*, *Gen. Rel. Grav.* **19** (1987) 1075-1082.
- [133] M. Gurses, *Perfect fluid sources in 2+1 dimensions*, *Class. Quantum Grav.* **11** (1994) 2585-2587.
- [134] A.K. Guts, *On the time-like closed smooth curves in general relativity*, *Izvestiya VUZov, Fizika*, n. **9** (1973) 33-36 (in Russian).

- [135] A.K. Guts, *Topological structure of the Gödel's universe*, *Izvestiya VUZov, Fizika*, n. **6** (1980) 109-110 (in Russian).
- [136] R. Hakim, *The inflationary universe: a primer*, in: "Gravitation, Geometry, and Relativistic Physics", *Lect. Notes Phys.* **212** (1984) 302-332.
- [137] D. Harari and P. Sikivie, *Effects of a Nambu-Goldstone boson on the polarization of radio galaxies and the cosmic microwave background*, *Phys. Lett.* **B289** (1992) 67-72.
- [138] E.R. Harrison, *On the origin of galactic rotation*, *Mon. Not. R. Astr. Soc.* **154** (1971) 167-186.
- [139] E.R. Harrison, *Galaxy formation and the early universe*, in: *Cargèse Lect. in Phys.* vol. **6**, Ed. E. Schatzman (Gordon and Breach: NY, 1973) 581-641.
- [140] W. Hasse, *The apparent size of distant objects*, *Gen. Rel. Grav.* **19** (1987) 515-524.
- [141] W. Hasse and V. Perlick, *Geometrical and kinematic characterization of parallax-free world models*, *J. Math. Phys.* **29** (1988) 2064-2069.
- [142] S.W. Hawking, *On the rotation of the universe*, *Mon. Not. R. Astr. Soc.* **142** (1969) 129-141.
- [143] S.W. Hawking and G.F.R. Ellis, *The large-scale structure of space-time* (Cambridge Univ. Press: Cambridge, 1973).
- [144] O. Heckmann and E. Schücking, *Ein Weltmodell der Newtonschen Kosmologie mit Expansion und Rotation*, in: "Jubilee of relativity theory", *Proc. Int. Conf., Bern, 11-17 July 1955*, Eds. A. Mercier and M. Kervaire, *Helv. Phys. Acta Suppl.* **IV** (1956) 114-115.
- [145] O. Heckmann and E. Schücking, *Newtonsche und Einsteinsche Kosmologie*, in: *Handbuch der Physik*, vol. *LIII*, Ed. S. Flügge (Springer: Berlin, 1959) 489-519.
- [146] O. Heckmann and E. Schücking, *Andere kosmologische Theorien*, in: *Handbuch der*

- Physik*, vol. *LIII*, Ed. S. Flügge (Springer: Berlin, 1959) 520-537.
- [147] O. Heckmann and E. Schücking, *Relativistic cosmology*, in: *Gravitation: An introduction to current research*, Ed. L. Witten (John Wiley: New York, 1962) 438-469.
 - [148] F.W. Hehl, P. von der Heyde, G.D. Kerlick, and J.M. Nester, *General relativity with spin and torsion: foundation and prospects*, *Revs. Mod. Phys.* **48** (1976) 393-416.
 - [149] F.W. Hehl, J.D. McCrea, E.W. Mielke, and Y. Ne'eman, *Metric-affine gauge theory of gravity: field equations, Noether identities, world spinors, and breaking of dilation invariance*, *Phys. Repts.* **258** (1995) 1-171.
 - [150] M. Heller, *Rotating bodies in general relativity*, *Acta Cosmologica* **3** (1975) 97-107.
 - [151] C.T. Hill, P.J. Steinhardt, and M.S. Turner, *Can oscillating physics explain an apparently periodic universe?*, *Phys. Lett.* **B252** (1990) 343-348.
 - [152] W.A. Hiscock, *Scalar perturbations in the Gödel universe*, *Phys. Rev.* **D17** (1978) 1497-1500.
 - [153] C. Hoenselaers and C.V. Vishveshwara, *A relativistically rotating fluid cylinder*, *Gen. Rel. Grav.* **10** (1979) 43-51.
 - [154] W.-H. Huang, *One-loop effective action on rotational spacetime – Zeta-function regularization and Schwinger perturbative expansion*, *Ann. Phys. (USA)* **254** (1997) 69-82.
 - [155] W.-H. Huang, *Chronology protection in generalized Gödel spacetime*, *Phys. Rev.* **D60** (1999) 067505 (3 pages).
 - [156] D. Hutsemekers, *Evidence for very large-scale coherent orientations of quasar polarization vectors*, *Astron. and Astrophys.* **332** (1998) 410-428.
 - [157] J. Hwang and H. Noh, *Cosmological vorticity in a gravity with quadratic order curvature couplings*, *Phys. Rev.* **D57** (1998) 2617-2620.

- [158] J.N. Islam, *Rotating fields in general relativity* (Cambridge Univ. Press: Cambridge, 1985) 122 p.
- [159] D.D. Ivanenko, *Rotation of the universe*, *Astron. Circ. Acad. Sci. USSR* **1254** (1983) 1-3 (in Russian).
- [160] Ivanenko D.D., & Krechet V.G., *Rotation of universe*, *Izvestia VUZov, Fizika*, n **3** (1986) 12-16 (in Russian).
- [161] D.D. Ivanenko, V.G. Krechet, and V.F. Panov, *On rotation of the universe*, in: “*Problems of gravity theory and elementary particles*”, Ed. K.P. Stanyukovich (Energoatomizdat: Moscow, 1986) v. **17**, p. 8-14 (in Russian).
- [162] D.D. Ivanenko, V.G. Krechet, and V.F. Panov, *Rotation of the universe and cosmology*, in: “*Gravitation and theory of relativity*”, Ed. V. R. Kajgorodov (Kazan Univ. Press: Kazan, 1987) v. **24**, p. 33-37 (in Russian).
- [163] D.D. Ivanenko and V.F. Panov, *Rotation of the universe and cosmology*, in: “*Problems of gravitation*”, Ed. D.V. Galtsov (Moscow Univ. Publ. House: Moscow, 1986) p. 168-175 (in Russian).
- [164] D.D. Ivanenko, V.A. Korotky, and Yu.N. Obukhov, *On the rotation of the universe*, *Astron. Circ. Acad. Sci. USSR* **1458** (1986) 1-3 (in Russian).
- [165] D.D. Ivanenko, V.A. Korotky, and Yu.N. Obukhov, *Cosmological scenario of a rotating universe*, *Astron. Circ. Acad. Sci. USSR* **1473** (1986) 1-3 (in Russian).
- [166] D.D. Ivanenko, V.A. Korotky, and Yu.N. Obukhov, *Microwave background radiation in a rotating and expanding universe*, *Astron. Circ. Acad. Sci. USSR* **1510** (1986) 2-4 (in Russian).
- [167] D.D. Ivanenko and Yu.N. Obukhov, *Model of the universe with expansion and rotation*, in: “*Prospects of unified theories*” eds. D.V. Galtsov, L.S. Kuzmenkov, and P.I. Pronin

- (Moscow Univ. Publ. House: Moscow, 1991) 98-115 (in Russian).
- [168] D.D. Ivanenko, P.I. Pronin, and G.A. Sardanashvily, *Gauge theory of gravity* (Moscow University Publ. House: Moscow, 1985), in Russian.
 - [169] P. Jain and J.P. Ralston, *Anisotropy in the distribution of galactic radio polarizations*, Preprint KUHEP-98-07 (Dept. Phys. Astr., Kansas Univ., Lawrence: 1998) 10 p.; e-print Archive: astro-ph/9803164.
 - [170] P. Jain and J.P. Ralston, *Anisotropy in the propagation of radio polarizations from cosmologically distant galaxies*, *Mod. Phys. Lett.*, **A14** (1999) 417-432.
 - [171] R. Jantzen, *Rotation in cosmology: comments on "Imparting rotation to a Bianchi type II space-time" by M.J. Rebouças and J.B.S. d'Oliveira [J. Math. Phys. 27, 417 (1986)] and similar papers*, *J. Math. Phys.* **27** (1986) 2748-2750.
 - [172] P. Kanti and C.G. Vayonakis, *Gödel-type universes in string-inspired charged gravity*, *Phys. Rev.* **D60** (1999) 103519 (13 pages).
 - [173] D.G. Kendall and G.A. Young, *Indirectional statistics and the significance of the asymmetric discovered by Birch*, *Mon. Not. R. Astr. Soc.* **207** (1984) 637-647.
 - [174] D.H. King, *A closed universe cannot rotate*, in: *Einstein Studies*, vol. 6: "Mach's principle: From Newton's bucket to quantum gravity", Eds. J.B. Barbour and H. Pfister (Birkhäuser: Boston, 1995), 237-248.
 - [175] A.R. King and G.F.R. Ellis, *Tilted cosmological models*, *Commun. Math. Phys.* **31** (1973) 209-242.
 - [176] C. Klein, *Rotational perturbations and frame dragging in a Friedman universe*, *Class. Quantum Grav.* **10** (1993) 1619-1631.
 - [177] C. Klein, *2nd-order effects of rotational perturbations of a Friedman universe*, *Class. Quantum Grav.* **11** (1994) 1539-1553.

- [178] P. Klepáč and J. Horský, *Charged universes of the Gödel type with closed timelike curves*, *Class. Quantum Grav.* **17** (2000) 2547-2555.
- [179] I.Yu. Kobzarev and K.G. Selivanov, *Rotation of the optical polarization vector in a nonstationary gravitational field*, *Sov. Phys. JETP* **67** (1988) 1955-1956 [*ZhETF* **94** n. 10 (1988) 1-4 (in Russian)].
- [180] A. Kogut, G. Hinshaw, and A.J. Banday, *Limits to global rotation and shear from the COBE DMR four year sky maps*, *Phys. Rev.* **D55** (1997) 1901-1905.
- [181] M.S. Kojima, *Slowly rotating charged viscous-fluid universe: exact solutions*, *Astrophys. Space Sci.* **143** (1988) 137-152.
- [182] M. Kokus, *Spherical rotation, particles and cosmology*, *Astrophys. and Space Sci.* **244** (1996) 353-356.
- [183] V.A. Korotky, *Primordial nucleosynthesis in the rotating and expanding universe*, in: "Modern Problems of Theoretical Physics. Festschrift for Professor D.Ivanenko", Eds. P.I. Pronin and Yu.N. Obukhov (World Scientific: Singapore, 1991) 121-127.
- [184] V.A. Korotky and V.G. Krechet, *Nonstationary cosmological models with rotation*, *Izvestiya VUZov, Fizika*, n. **3** (1988) 48-51 (in Russian).
- [185] V.A. Korotky and V.G. Krechet, *Self-consistent solutions in cosmological models with rotation*, *Izvestiya VUZov, Fizika*, n. **6** (1988) 5-10 (in Russian).
- [186] V.A. Korotky and Yu.N. Obukhov, *Microwave background radiation in rotating Universe*, *Preprint IFT/22/87* (Inst. Theor. Phys., Warsaw Univ.: Warsaw, 1987) 11 p.
- [187] V.A. Korotky and Yu.N. Obukhov, *Kinematical analysis of cosmological models with rotation*, *Sov. Phys. JETP* **72**(1) (1991) 11-15 [*ZhETF* **99** 1 (1991) 22-31 (in Russian)].
- [188] V.A. Korotky and Yu.N. Obukhov, *Electromagnetic waves in the rotating Universe*, *Mosc. Univ. Phys. Bull.* **46**, n. **2** (1991) 4-6 [*Vestnik Mosk. Univ., ser. fizika*, as-

- tronomiya, **32**, n. 2 (1991) 3-6 (in Russian)].
- [189] V.A. Korotky and Yu.N. Obukhov, *Rotating and expanding cosmology in ECSK-theory*, *Astrophys. Space Sci.* **198** (1992) 1-12.
- [190] V.A. Korotky and Yu.N. Obukhov, *General-relativistic cosmology with expansion and rotation*, *Russ. Phys. J.* **36**, n **6** (1993) 568-573 [*Izvestiya VUZov, Fizika*, n. **6** (1993) 71-77 (in Russian)].
- [191] V.A. Korotky and Yu.N. Obukhov, *Bianchi IX cosmological models with rotation in general relativity theory*, *Russ. Phys. J.* **37**, n **9** (1994) 909-914 [*Izvestiya VUZov, Fizika*, n. **9** (1994) 122-127 (in Russian)].
- [192] V.A. Korotky and Yu.N. Obukhov, *Bianchi II cosmological models with rotation in general relativity theory*, *Russ. Phys. J.* **37**, n **10** (1994) 961-970 [*Izvestiya VUZov, Fizika*, n. **10** (1994) 54-63 (in Russian)].
- [193] V.A. Korotky and Yu.N. Obukhov, *Can cosmic rotation explain an apparently periodic universe?*, *Gen. Relat. Grav.* **26** (1994) 429-435.
- [194] V.A. Korotky and Yu.N. Obukhov, *Polarization of radiation in a rotating universe*, *JETP* **81** (6) (1995) 1031-1035 [*ZhETF* **108** 6(12) (1995) 1889-1898 (in Russian)].
- [195] V.A. Korotky and Yu.N. Obukhov, *On cosmic rotation*, in: "Gravity, Particles and Space-Time" /Eds. P.Pronin and G.Sardanashvily (World Scientific: Singapore, 1996) 421-439.
- [196] V.A. Korotky and Yu.N. Obukhov, *Bianchi-II rotating world*, *Astrophys. Space Sci.* **260** (1999) 425-439.
- [197] D. Kramer, H. Stephani, M. MacCallum, E. Herlt, *Exact solutions of the Einstein's field equations*, Ed. E. Schmutzer (Deutscher Verlag der Wissenschaften: Berlin, 1980).
- [198] A. Kasiński, *Solutions of the Einstein field equations for a rotating perfect fluid. Part*

1. *Presentation of the flow-stationary and vortex-homogeneous solutions*, *Acta Phys. Pol.* **B5** (1974) 411-436.
- [199] A. Krasinski, *Solutions of the Einstein field equations for a rotating perfect fluid. Part 2. Properties of the flow-stationary and vortex-homogeneous solutions*, *Acta Phys. Pol.* **B6** (1975) 223-238.
- [200] A. Krasinski, *Solutions of the Einstein field equations for a rotating perfect fluid. Part 3. Survey of models of rotating perfect fluid or dust*, *Acta Phys. Pol.* **B6** (1975) 239-240.
- [201] A. Krasinski, *Some solutions of the Einstein field equations for a rotating perfect fluid*, *J. Math. Phys.* **16** (1975) 125-131.
- [202] A. Krasinski, *All flow-stationary cylindrically symmetric solutions of the Einstein field equations for a rotating isentropic perfect fluid*, *Rept. Math. Phys.* **14** (1978) 225-236.
- [203] A. Krasinski, *Rotational motion of matter in general relativity*, *Acta Cosmologica* **7** (1978) 119-131.
- [204] A. Krasinski, *Cylindrical rotating universe*, *Acta Cosmologica* **7** (1978) 133-137.
- [205] A. Krasinski, *Bibliography on inhomogeneous cosmological models*, *Acta Cosmologica* **20** (1994) 67-98.
- [206] A. Krasinski, *Rotating dust solutions of Einstein's equations with 3dimensional symmetry groups. Part 1: Two Killing fields spanned on u^α and w^α* , *J. Math. Phys.* **39** (1998) 380-400.
- [207] A. Krasinski, *Rotating dust solutions of Einstein's equations with three-dimensional symmetry groups. Part 2: One Killing field spanned on u^α and w^α* , *J. Math. Phys.* **39** (1998) 401-422.
- [208] A. Krasinski, *Rotating dust solutions of Einstein's equations with three-dimensional symmetry groups. Part 3: All Killing fields linearly independent of u^α and w^α* , *J.*

- Math. Phys.* **39** (1998) 2148-2179.
- [209] A. Krasinski, *Inhomogeneous cosmological models* (Cambridge Univ. Press: Cambridge, 1997) 317 pp.
- [210] A. Krasinski, *The ultimate extension of the Bianchi classification for rotating dust models*, in: “On Einstein’s path”, *Essays in honor of E. Schucking*, Ed. A. Harvey (Springer: New York, 1998) 283-298.
- [211] V.G. Krechet, *Rotating quantum cosmological models with wave fields*, *Russ. Phys. J.* **36**, n. 1 (1993) 33-35 [*Izvestiya VUZov, Fizika* **n. 1** (1993) 41-44 (in Russian)].
- [212] V.G. Krechet and V.F. Panov, *Nonstationary cosmological models with rotation*, *Astrophysics (USA)* **28** (1988) 400-405 [*Astrofizika* **28** (1988) 670-678 (in Russian)].
- [213] V.G. Krechet and I.V. Sandina, *Self-gravitating wave fields in Gödel space*, *Izvestiya VUZov, Fizika* **n. 10** (1981) 59-63 (in Russian).
- [214] J. Kristian and R. Sachs, *Observations in cosmology*, *Astrophys. J.* **143** (1966) 379-399.
- [215] K.D. Krori, P. Borgohain, P.K. Kar, and Dipali Das (Kar), *Exact scalar and spinor solutions in some rotating universes*, *J. Math. Phys.* **29** (1988) 1645-1649.
- [216] R.W. Kühne, *On the cosmic rotation axis*, *Mod. Phys. Lett.* **A12** (1997) 2473-2474.
- [217] W. Kundt, *Trägheitsbahnen in einem von Gödel angegebenen kosmologischen Modell*, *Z. Physik* **145** (1956) 611-620.
- [218] E.V. Kuvshinova and V.F. Panov, *5-dimensional nonstationary cosmological models with rotation*, *Izvestiya VUZov, Fizika*, n. **9** (1997) 25-28 (in Russian).
- [219] A.A. Kurskov and L.M. Ozernoi, *Evolution of cosmological turbulence. I*, *Sov. Astron.* **18** (1974) 157 [*Astron. Zh.* **51** (1974) 270 (in Russian)].
- [220] A.A. Kurskov and L.M. Ozernoi, *Evolution of cosmological turbulence. II. Dissipa-*

- tion of subsonic motions and alternative modes of post-recombination evolution*, *Sov. Astron.* **18** (1974) 300-306 [*Astron. Zh.* **51** (1974) 508-519 (in Russian)].
- [221] A.A. Kurskov and L.M. Ozernoi, *Evolution of cosmological turbulence. III. Spectrum of density inhomogeneities produced by turbulence*, *Sov. Astron.* **18** (1974) 700-707 [*Astron. Zh.* **51** (1974) 1177-1190 (in Russian)].
- [222] A.A. Kurskov and L.M. Ozernoi, *Evolution of cosmological turbulence. IV. Gravitational growth of inhomogeneities with isolation of galaxies and galaxy systems*, *Sov. Astron.* **19** (1976) 569-576 [*Astron. Zh.* **52** (1975) 937-949 (in Russian)].
- [223] A.A. Kurskov and L.M. Ozernoi, *Angular fluctuations of relict radiation produced by cosmological turbulence*, *Astrophys. and Space Sci.* **56** (1978) 67-80.
- [224] Z. Lalak, S. Pokorski, and J. Wess, *Spin-1/2 particle in gravitational field of a rotating body*, *Phys. Lett.* **B355** (1995) 453-460.
- [225] K. Lanczos, *Über eine stationäre Kosmologie im Sinne der Einsteinschen Gravitationstheorie*, *Zeits. f. Phys.* **21** (1924) 73-110 [English translation: *On a Stationary cosmology in the sense of Einstein's theory of gravitation*, *Gen. Rel. Grav.* **29** (1997) 361-399].
- [226] B.E. Laurent, K. Rosquist, and E. Sviestins, *The behavior of null geodesics in a class of rotating space-time homogeneous cosmologies*, *Gen. Rel. Grav.* **13** (1981) 1093-1116.
- [227] J. Lathrop and R. Teglas, *Dynamics in the Gödel universe*, *Nuovo Cim.* **B43** (1978) 162-172.
- [228] D. Leahy, *Scalar and neutrino fields in the Gödel universe*, *Int. J. Theor. Phys.* **21** (1982) 703-753.
- [229] J.P. Leahy, *Comment on the measurement of cosmological birefringence*, Preprint University of Manchester (1997) 1 p. (e-Print Archive: astro-ph/9704285).

- [230] G. Lemaître, *L'hypothèse de l'atome primitif*, *Acta Pontifica Acad. Sci.* **12** (1948) 25-40.
- [231] T. Lewis, *Some special solutions of the equations of axially symmetric gravitational fields*, *Proc. Roy. Soc. Lond.* **A136** (1932) 176-192.
- [232] Li-Xin Li, *Effect of the global rotation of the universe on the formation of galaxies*, *Gen. Relat. Grav.* **30** (1998) 497-507.
- [233] A. Linde, *Particle physics and inflationary cosmology* (Harwood: Chur, 1990).
- [234] T.J. Loredo, E.E. Flanagan, and I.M. Wasserman, *Bayesian analysis of the polarization of distant radio sources: Limits on cosmological birefringence*, *Phys. Rev.* **D56** (1997) 7507-7512.
- [235] D. Lorenz-Petzold, *Electromagnetic rotating universes*, *Astrophys. and Space Sci.* **96** (1983) 343-349.
- [236] V.N. Lukash, I.D. Novikov, and A.A. Starobinsky, *Particle creation in the vortex cosmological model*, *Sov. Phys. JETP* **42** (1976) 757-765 [*ZhETF* **69**, no. 11 (1975) 1484-1500 (in Russian)].
- [237] R. Maartens, G.F.R. Ellis, and W.R. Stoeger, *Anisotropy and inhomogeneity of the universe from $\Delta T/T$* , *Astron. and Astrophys.* **309** (1996) L7-L10.
- [238] M.A.H. MacCallum, *Inhomogeneous and anisotropic cosmologies*, in: "The origin of structure in the universe" (NATO ASI Series C, vol. 393) Eds. E. Gunzig and P. Nardone (Kluwer: Dordrecht, 1993) 131-159.
- [239] M.A.H. MacCallum, *Anisotropic and inhomogeneous cosmologies*, in: *The renaissance of general relativity and cosmology (Festschrift for D. Sciama)*, Eds. G.F.R. Ellis, A. Lanza, and J. Miller (Cambridge Univ. Press: Cambridge, 1993) 213-233.
- [240] M.A.H. MacCallum and G.F.R. Ellis, *A class of homogeneous cosmological models. II.*

- Observations, Commun. Math. Phys.* **19** (1970) 31-64.
- [241] M.N. Macrossan, *A thermodynamic origin of a universal mass-angular momentum relationship*, *Astrophys. and Space Sci.* **133** (1987) 403-410.
- [242] S. Maitra, *Stationary dust-filled cosmological solution with $\Lambda = 0$ and without closed timelike lines*, *J. Math. Phys.* **7** (1966) 1025-1030.
- [243] A.V. Mandzhos and V.V. Tel'nyuk-Adamchuk, *On the preference orientation of galaxy clusters*, *Astron. Circ. Acad. Sci. USSR* **681** (1972) 1-2 (in Russian).
- [244] A.V. Mandzhos and V.V. Tel'nyuk-Adamchuk, *The galaxy clusters orientation and the Metagalaxy anisotropy*, *Astron. Circ. Acad. Sci. USSR* **767** (1973) 1-3 (in Russian).
- [245] A.V. Mandzhos and V.V. Tel'nyuk-Adamchuk, *On the orientations of the Uppsala catalogue's galaxies*, *Astron. Circ. Acad. Sci. USSR* **908** (1976) 5-8 (in Russian).
- [246] K. Maniharsingh, *Rotational perturbations of cosmological viscous-fluid universe with zero-mass scalar field*, *Astrophys. and Space Sci.* **139** (1987) 183-193.
- [247] K. Maniharsingh, *Slowly-rotating charged viscous-fluid universes – Exact solutions*, *Astrophys. and Space Sci.* **143** (1988) 137-152.
- [248] K. Maniharsingh, *Slow rotation of radiating viscous-fluid universe coupled with scalar field*, *Astrophys. and Space Sci.* **148** (1988) 149-161.
- [249] K. Maniharsingh, *Slowly rotating perfect-fluid universes coupled with zero-mass scalar field*, *Astrophys. and Space Sci.* **161** (1989) 111-123.
- [250] K. Maniharsingh, *Rotational perturbations of charged cosmological viscous-fluid universes coupled with a scalar field*, *Astrophys. and Space Sci.* **162** (1989) 29-150.
- [251] K. Maniharsingh, *Rotating magnetoviscous-fluid universes in general relativity*, *Astrophys. and Space Sci.* **164** (1990) 153-166.

- [252] K. Maniharsingh, *Rotating charged perfect-fluid universes coupled with scalar field in general relativity*, *Astrophys. and Space Sci.* **172** (1990) 29-45.
- [253] K. Maniharsingh, *Rotational perturbations of magneto-viscous fluid universes coupled with zero-mass scalar field*, *Astrophys. and Space Sci.* **180** (1991) 253-271.
- [254] K. Maniharsingh, *Particle creation in slowly-rotating Robertson-Walker universes*, *Astrophys. and Space Sci.* **182** (1991) 141-153.
- [255] K. Maniharsingh, *On slowly rotating cosmological viscous fluid model universes in general relativity*, *Astrophys. and Space Sci.* **182** (1991) 207.
- [256] K. Maniharsingh, *Rotational perturbations of relativistic perfect-fluid model universes interacting with gravitational field*, *Astrophys. and Space Sci.* **183** (1991) 285-307.
- [257] R. Mansouri and K. Nouzari, *Rotation of plane of polarization of an electromagnetic wave propagating over cosmological distances in "Finite rotating universe" scenario*, *Preprint Institute for Studies in Physics and Mathematics, Tehran, Iran* (1997) 4 p.; e-print Archive: astro-ph/9710028.
- [258] B. Mashhoon, *Influence of gravitation on the propagation of electromagnetic radiation*, *Phys. Rev.* **D11** (1975) 2679-2684.
- [259] B. Mashhoon, *On the relativity of rotation*, in: *Directions in general relativity (Festschrift for D.Brill)*, vol. 2, Eds. B.L. Hu and T.A. Jacobson (Cambridge Univ. Press: Cambridge, 1993) 182-194.
- [260] D.R. Matravers, M.D. Madsen, and D.L. Vogel, *The microwave background and (m, Z) relations in a tilted cosmological model*, *Astrophys. and Space Sci.* **112** (1985) 193-202.
- [261] R.A. Matzner, *Closed rotating cosmologies containing matter described by the kinetic theory. A: Formalism*, *Ann. Phys.* **65** (1971) 438-481.
- [262] R.A. Matzner, *Closed rotating cosmologies containing matter described by the kinetic*

- theory. B: Small anisotropy; application to observations*, *Ann. Phys.* **65** (1971) 482-505.
- [263] R.A. Matzner and B.W. Tolman, *Cosmological spatial curvature probed by microwave polarization*, *Phys. Rev.* **D26** (1982) 2957-2964.
- [264] R.A. Matzner, T. Rothman, and G.F.R. Ellis, *Conjecture on isotope production in the Bianchi cosmologies*, *Phys. Rev.* **D34** (1986) 2926-2933.
- [265] S. Mavrides, *L'Univers relativiste* (Masson: Paris, 1973), p. 178-180.
- [266] E.W. Mielke, *Geometrodynamics of Gauge Fields — On the geometry of Yang–Mills and gravitational gauge theories* (Akademie–Verlag: Berlin, 1987).
- [267] Hokee Minn, *Global theory of Einstein–Cartan equations: Gödel universe with torsion*, *J. Korean Astron. Soc.* **9** (1976) 9-14.
- [268] W. Monteiro S.Jr., *Stably causal Gödel-type models*, *J. Math. Phys.* **32** (1991) 3223-3227.
- [269] W. Monteiro S.Jr. and N.A. Tomimura, *Existence and causality of cylindrically symmetric cosmological models with rotating spin fluids*, *Class. Quantum Grav.* **8** (1991) 977-984.
- [270] W. Monteiro S.Jr. and N.A. Tomimura, *Existence and causality of cosmological models with spinning fluids and torsion*, *Phys. Lett.* **A197** (1995) 267-274.
- [271] R.M. Muradian, *On the origin of rotation of galaxies in cosmology of Ambartsumyan*, *Astrofizika* **11** (1975) 237-248 (in Russian).
- [272] R.M. Muradian, *Cosmic numbers and rotation of the metagalaxy*, *Astrofizika* **13** (1977) 63-67 (in Russian).
- [273] R.M. Muradian, *The primeval hadron: origin of stars, galaxies and astronomical universe*, *Astrophys. and Space Sci.* **69** (1980) 339-351 [*Astrophys. and Space Sci.* **69** (1980) 325-337 (in Russian)].

- [274] R.M. Muradian, *Cosmological constant and rotation of the universe*, *Astrofizika* **21** (1984) 396.
- [275] R.M. Muradian, S. Carneiro, and R. Marques, *Radius-mass scaling laws for celestial bodies*, e-print Archive: astro-ph/9907129.
- [276] H. Nariai, *Generation of sound from primordial cosmic turbulence*, in: “Confrontation of cosmological theories with observational data”, *Proc. of the IAU Symp. N. 63, Cracow, Poland, September 10-12, 1973*, Ed. M.S. Longair (D. Reidel Publishing Co.: Dordrecht, 1974) 241-243.
- [277] J.V. Narlikar, *Rotating universes*, in: “Evidence for gravitational theories”, *Proc. of Int. School of Physics “E.Fermi”, Course XX, Varenna, June 19-July 1, 1961*, Ed. C. Moller (Academic Press: NY, 1962) 222-227.
- [278] J.V. Narlikar, *Spinning Universes in Newtonian cosmology*, in: “On Einstein’s path”, *Essays in honor of E. Schucking*, Ed. A. Harvey (Springer: New York, 1998) 319-327.
- [279] B. Nodland and J.P. Ralston, *Indication of anisotropy in electromagnetic propagation over cosmological distances*, *Phys. Rev. Lett.* **78** (1997) 3043-3046.
- [280] B. Nodland and J.P. Ralston, *Appropriate null hypothesis for cosmological birefringence - Reply*, *Phys. Rev. Lett.* **79** (1997) 1958.
- [281] B. Nodland and J.P. Ralston, *Response to Leahy’s comment on the data’s indication of cosmological birefringence*, *Preprint Univ. of Rochester* (1997) 3 p.; e-print Archive: astro-ph/9706126.
- [282] B. Nodland and J.P. Ralston, *Cosmologically screwy light*, in: “Particles and nuclei”, *Proc. of the 14th Particles and Nuclei International Conference (PANIC96)*, Williamsburg, Virginia, May 22-28, 1996. Eds. C. Carlson and John J. Domingo (World Scientific: Singapore, 1997) 485-486.

- [283] M. Novello and M. Rebouças, *The stability of a rotating universe*, *Astrophys. J.* **225** (1978) 719-724.
- [284] M. Novello and M. Rebouças, *Rotating universe with successive causal and non-causal regions*, *Phys. Rev.* **D19** (1979) 2850-2852.
- [285] M. Novello, I. Damião Soares, and J. Tiomno, *Geodesic motion and confinement in Gödel's universe*, *Phys. Rev.* **D27** (1983) 779-788.
- [286] I.D. Novikov, V.P. Frolov, *Physics of black holes* (Nauka: Moscow, 1986), in Russian.
- [287] Yu.N. Obukhov, *Observations in rotating cosmologies*, in: "Gauge Theories of Fundamental Interactions" (Proceedings of the XXXII Semester in Stefan Banach Int. Math. Center, Warsaw, Poland, 1988)/Eds. R. Rączka and M. Pawłowski (World Scientific: Singapore, 1990) 341-366.
- [288] Yu.N. Obukhov, *Rotation in cosmology*, *Gen. Relat. Grav.* **24** (1992) 121-128.
- [289] Yu.N. Obukhov and V.A. Korotky, *The Weyssenhoff fluid in Einstein-Cartan theory*, *Class. Quantum Grav.* **4** (1987) 1633-1657.
- [290] Yu.N. Obukhov and O.B. Piskareva, *Spinning fluid in general relativity*, *Class. Quantum Grav.* **6** (1989) L15-L19.
- [291] Yu.N. Obukhov and O.B. Piskareva, *The development of small perturbations of cosmological models in the general theory of relativity: spin effects*, *Sov. Phys. JETP* **71**(1) (1990) 1-11 [*ZhETF* **98** 1(7) (1990) 3-24 (in Russian)].
- [292] Yu.N. Obukhov, V.A. Korotky, and F.W. Hehl, *On the rotation of the universe*, *Preprint Univ. of Cologne* (1997) 1 p.; e-print Archive: astro-ph/9705243.
- [293] J.H. Oort, *The formation of galaxies and the origin of the high-velocity hydrogen*, *Astron. and Astrophys.* **7** (1970) 381-404.
- [294] L.M. Ozernoy, *Whirl theory of the origin of galaxies and clusters of galaxies*, in: "Con-

- frontation of cosmological theories with observational data”, *Proc. of the IAU Symp. N. 63, Cracow, Poland, September 10-12, 1973*, Ed. M.S. Longair (D. Reidel Publishing Co.: Dordrecht, 1974) 227-240.
- [295] I. Ozsváth and E. Schücking, *Finite rotating universe*, *Nature* **193** (1962) 1168-1169.
- [296] I. Ozsváth and E. Schücking, *An anti-Mach metric*, in: *Recent developments in general relativity* (Pergamon: Oxford, 1962) 339-350.
- [297] I. Ozsváth and E. Schücking, *The finite rotating universe*, *Ann. Phys.* **55** (1969) 166-204.
- [298] I. Ozsváth and E.L. Schücking, *The world viewed from outside*, *J. Geom. Phys.* **24** (1998) 303-333.
- [299] I. Ozsváth, *The finite rotating universe revisited*, *Class. Quantum Grav.* **14** (1997) A291-A297.
- [300] I. Ozsváth, *New homogeneous solutions of Einstein’s field equations with incoherent matter obtained by a spinor technique*, *J. Math. Phys.* **6** (1965) 590-610.
- [301] I. Ozsváth, *Two rotating universe with dust and electromagnetic field*, in: *Perspectives in Geometry and Relativity*, Ed. B. Hoffmann (Indiana Univ. Press: Bloomington, 1966) 245-258.
- [302] I. Ozsváth, *Spatially homogeneous world models*, *J. Math. Phys.* **11** (1970) 2860-2870.
- [303] I. Ozsváth, *Dust-filled universes of class II and class III*, *J. Math. Phys.* **11** (1970) 2871-2883.
- [304] I. Ozsváth, *Spatially homogeneous rotating world models*, *J. Math. Phys.* **12** (1971) 1078-1082.
- [305] I. Ozsváth, *Working with Engelbert*, in: “On Einstein’s path”, *Essays in honor of E. Schucking*, Ed. A. Harvey (Springer: New York, 1998) 339-351.

- [306] J. Pachner, *Rotating incoherent matter in general relativity*, *Can. J. Phys.* **51** (1973) 477-490.
- [307] L.A. Page, *On observing the cosmic microwave background*, in: *Proc. of NATO Adv. Study Inst.: 3rd Erice School on Particle Astrophysics: Generation of Large Scale Cosmological Structures, Erice, Italy, 3-13 Nov 1996* (e-Print Archive: astro-ph/9703054) 30 pp.
- [308] D. Palle, *Imperfect fluid in the Einstein-Cartan gravity*, Preprint Dept. Theor. Phys., R.Boskovic Inst. (July 1995) 6 p.
- [309] D. Palle, *On certain relationships between cosmological observables in the Einstein-Cartan gravity*, *Nuovo Cim.* **B111** (1996) 671-675.
- [310] D. Palle, *On the area distance and the Riemannian geometry*, Preprint Dept. Theor. Phys., R.Boskovic Inst. (July 1999) 5 p; e-print Archive: astro-ph/9905252.
- [311] V.F. Panov, *Study of the rotation of the universe*, *Izvestiya VUZov, Fizika*, n. **1** (1985) 22-25 (in Russian).
- [312] V.F. Panov, *Rotation of the early universe*, *Sov. Phys. J.* **28** (1985) 981-983 [*Izvestiya VUZov, Fizika*, n. **12** (1985) 37-40 (in Russian)].
- [313] V.F. Panov, *Problem of the universe's rotation*, *Izvestiya VUZov, Fizika*, n. **9** (1987) 58-62 (in Russian).
- [314] V.F. Panov, *Spontaneous symmetry breaking in cosmological models with rotation*, *Teor. Mat. Fiz.* **74** (1988) 463-468 (in Russian).
- [315] V.F. Panov, *Bianchi type VIII rotating cosmological models*, *Izvestiya VUZov, Fizika*, n. **5** (1989) 98-103 (in Russian).
- [316] V.F. Panov, *Rotating models of the early universe*, *Izvestiya VUZov, Fizika*, n. **6** (1989) 67-70 (in Russian).

- [317] V.F. Panov, *Spontaneous breaking of the T-symmetry in a Gödel type cosmological model*, *Izvestiya VUZov, Fizika*, n. **9** (1989) 14-16 (in Russian).
- [318] V.F. Panov, *Nonstationary Gödel type cosmological model*, *Izvestiya VUZov, Fizika*, n. **1** (1990) 62-66 (in Russian).
- [319] V.F. Panov, *A cosmological model with rotation in the relativistic theory of gravity*, *Theor. and Math. Phys.* **114** (1998) 249-253.
- [320] V.F. Panov, *Scalar field in a nonstationary Gödel type cosmological model*, *Izvestiya VUZov, Fizika*, n. **2** (1991) 54-57 (in Russian).
- [321] V.F. Panov and Yu.G. Sbytov, *On the possibility to explain the observational Birch anisotropy with the universe rotation*, *Sov. Phys. JETP* **74** (1992) 411-415 [*ZhETF* **101** (1992) 769-778 (in Russian)].
- [322] V.F. Panov and Yu.G. Sbytov, *Behavior of a bundle of rays forming the image of a source in cosmological models with rotation*, *JETP* **87** (1998) 417-420 [*ZhETF* **114** (1998) 769-776 (in Russian)].
- [323] S.L. Parnovskii, Yu.N. Kudrya, and A.N. Aleksandrov, *Apparent anisotropy in the orientation of extragalactic objects due to the curvature of spacetime*, *JETP* **79** (1994) 840-846 [*ZhETF* **106** (1994) 1559-1571 (in Russian)].
- [324] L.K. Patel and P.C. Vaidya, *A note on rotating universe*, *Current Sci. (India)* **38** (1969) 334.
- [325] L.K. Patel and N. Dadhich, *A class of stationary rotating string cosmological models*, *Pramaña* **47** (1996) 387-392.
- [326] L.K. Patel and S.D. Maharaj, *Stationary rotating string world models with a magnetic field*, *Pramaña* **47** (1996) 33-39.
- [327] L.K. Patel and S.K. Koppar, *A rotating mass in Gödel universe with an electromagnetic*

- field*, *Acta Phys. Hungarica* **61** (1987) 363-367.
- [328] M.D. Patel and P.C. Vaidya, *A rotating mass embedded in an Einstein-Gödel universe*, *Gen. Rel. Grav.* **15** (1983) 777-783.
- [329] V.N. Pavelkin and V.F. Panov, *A study of rotation in cosmology*, *Izvestiya VUZov, Fizika*, n. **7** (1988) 29-32 (in Russian).
- [330] V.N. Pavelkin and V.F. Panov, *A nonstationary cosmological model with rotation in the Einstein-Cartan theory*, *Russ. Phys. J.* **36**, n **8** (1993) 784-788 [*Izvestiya VUZov, Fizika*, n. **8** (1993) 90-94 (in Russian)].
- [331] V.N. Pavelkin and V.F. Panov, *Large scale anisotropy of microwave background radiation in rotating cosmologies*, *Int. J. Mod. Phys.* **D4** (1995) 161-165.
- [332] P.R.C.T. Pereira, N.O. Santos, and A.Z. Wang, *Geodesic motion and confinement in Lanczos spacetime*, *Class. Quantum Grav.* **13** (1996) 1641-1654.
- [333] Z. Perjés, G. Fodor, L.A. Gergely, and M. Marklund, *A rotating incompressible perfect fluid space-time*, *Preprint KFKI Budapest* (1998) 3 p.; e-print Archive: gr-qc/9806095.
- [334] V. Perlick and W. Hasse, *Gravitational Faraday effect in conformally stationary spacetimes*, *Class. Quantum Grav.* **10** (1993) 147-161.
- [335] E.S. Phinney and R.L. Webster, *Is there evidence for universal rotation?*, *Nature* **301** (1982) 735-736.
- [336] J. Pfarr, *Time travel in Gödel's space*, *Gen. Relat. Grav.* **13** (1981) 1073-1091.
- [337] L.O. Pimentel and A. Macías, *Klein-Gordon and Weyl equations in the Gödel universe*, *Phys. Lett.* **A117** (1986) 325-327.
- [338] L.O. Pimentel, A. Camacho, and A. Macías, *Weyl equation in Gödel type universes*, *Mod. Phys. Lett.* **A9** (1994) 3707-3712.

- [339] T. Piran and P.N. Safier, *A gravitational analogue of Faraday rotation*, *Nature* **318** (1985) 271-273.
- [340] J. Plebanski, *Electromagnetic waves in gravitational fields*, *Phys. Rev.* **118** (1960) 1396-1408.
- [341] V.N. Ponomarev, A.O. Barvinsky, and Yu.N. Obukhov, *Geometrodynamical methods and the gauge approach to the theory of gravitational interactions* (Energoatomizdat: Moscow, 1985), in Russian.
- [342] D.J. Raine, *Mach's principle and space-time structure*, *Rep. Prog. Phys.* **44** (1981) 1151-1195.
- [343] E. Radu, *On the Euclidean approach to quantum field theory in Gödel space-time*, *Phys. Lett.* **A247** (1998) 207-210.
- [344] E. Radu, *Features of a charged rotating spacetime*, *Class. Quantum Grav.* **15** (1998) 2743-2760.
- [345] E. Radu, *On a Gödel-type Euclidean solution*, *Gen. Rel. Grav.* **31** (1999) 287-300.
- [346] E. Radu and D. Amariutei, *On a new Petrov type-I rotating solution*, *Nuovo Cim.* **B113** (1998) 1317-1321.
- [347] J.P. Ralston and B. Nodland, *An update on cosmological anisotropy in electromagnetic propagation*, in: "Big Sky 1997, Intersections between particle and nuclear physics", *Proc. of the 7th Int. Conf. on the Intersections of Particle and Nuclear Physics (Big Sky, Montana, 1997)*, Ed. T. W. Donnelly (American Institute of Physics, 1997) 432-437; e-print Archive: astro-ph/9708114.
- [348] J.P. Ralston, P. Jain and B. Nodland, *New phenomena in propagation of radio polarizations due to magnetic fields on cosmological scales*, *Phys. Rev. Lett.* **81** (1998) 26-29.

- [349] A. Ratnaprabha, and R.K. Tarachand Singh, *Cosmological viscous fluid universe in slow rotation*, *Astrophys. and Space Sci.* **165** (1990) 175-187.
- [350] D. Ray, *Gödel-like cosmological solutions*, *J. Math. Phys.* **21** (1980) 2797-2798.
- [351] A. Raychaudhuri, *Relativistic cosmology. I*, *Phys. Rev.* **98** (1955) 1123-1126.
- [352] A. Raychaudhuri, *Theoretical cosmology*, (Clarendon Press: Oxford, 1979) 371 p.
- [353] A. Raychaudhuri, *An approach to anisotropic cosmologies*, in: “Gravitation, Gauge Theories and the Early Universe”, Eds. B.R. Iyer, C.V. Vishveshwara, and N. Mukunda (Kluwer: Dordrecht, 1989) 89-106.
- [354] A. Raychaudhuri and S. Thakurta, *Homogeneous spacetimes of the Gödel type*, *Phys. Rev.* **D22** (1980) 802-806.
- [355] A. Rebhan, *Large-scale rotational perturbations of a Friedmann universe with collisionless matter and primordial magnetic fields*, *Astrophys. J* **392** (1992) 385-393.
- [356] M.J. Rebouças, *A rotating universe with violation of causality*, *Phys. Lett.* **A70** (1979) 161-163.
- [357] M.J. Rebouças, *Time-dependent, Bianchi II, rotating universe*, *Nuovo Cim.* **B67** (1982) 120-126.
- [358] M.J. Rebouças and J.E. Āman, *Computer aided study of a class of Riemannian spacetimes*, *J. Math. Phys.* **28** (1987) 888-896.
- [359] M.J. Rebouças and J.A.S. de Lima, *Time-dependent, finite, rotating universe*, *J. Math. Phys.* **22** (1981) 2699-2703; *Erratum*, *J. Math. Phys.* **23** (1982) 1547.
- [360] M.J. Rebouças and J. Tiomno, *Homogeneity of Riemannian spacetimes of Gödel type*, *Phys. Rev.* **D28** (1983) 1251-1264.
- [361] M.J. Rebouças and J. Tiomno, *A class of inhomogeneous Gödel-type models*, *Nuovo*

- Cim.* **B90** (1985) 204-210.
- [362] M.J. Rebouças and J.B.S. d’Olival, *Imparting rotation to a Bianchi type II space-time*, *J. Math. Phys.* **27** (1986) 417-418; *Erratum*, *J. Math. Phys.* **27** (1986) 2209.
- [363] M.J. Rebouças, J.E. Åman, and A.F.F. Teixeira, *A note on Gödel-type space-time*, *J. Math. Phys.* **27** (1986) 1370-1372.
- [364] M.J. Rebouças and A.F.F. Teixeira, *Features of a relativistic space-time with seven isometries*, *Phys. Rev.* **D34** (1986) 2985-2989.
- [365] M.J. Rebouças and A.F.F. Teixeira, *Riemannian space-times of Gödel type in five dimensions*, *J. Math. Phys.* **39** (1998) 2180-2192.
- [366] M.J. Rebouças and A.F.F. Teixeira, *Causal anomalies in Kaluza-Klein gravity theories*, *Int. J. Mod. Phys.* **13** (1998) 3181-3191.
- [367] W. Rindler and V. Perlick, *Rotating coordinates as tools for calculating circular geodesics and gyroscopic precession*, *Gen. Rel. Grav.* **22** (1990) 1067-1081.
- [368] M. Rooman and Ph. Spindel, *Gödel metric as a squashed anti-de Sitter geometry*, *Class. Quantum Grav.* **15** (1998) 3241-3249.
- [369] K. Rosquist, *Global rotation*, *Gen. Rel. Grav.* **12** (1980) 649-664.
- [370] K. Rosquist, *Exact rotating and expanding radiation-filled universe*, *Phys. Lett.* **A97** (1983) 145-146.
- [371] K. Rosquist, *Trigonometric parallaxes of distant objects: what they could tell about the universe*, *Astrophys. J.* **331** (1988) 648-652.
- [372] V.A. Ruban and A.D. Chernin, *Rotational perturbations of anisotropic cosmology*, *Astrofiz. Zhurn.* **49** (1972) 447-449 (in Russian).
- [373] G. Ruben, *Die Rotation des Kosmos aus der Aberration von Quasaren*, *Ann. d. Phys.*

- 44** (1987) 150-152.
- [374] G. Ruben, *Cosmic rotation and the inertial system*, *Astrophys. and Space Sci.* **177** (1991) 465-470.
- [375] T.V. Ruzmaikina and A.A. Ruzmaikin, *Evolution of a cosmological model with rotation*, *Sov. Phys. JETP.* **29** (1969) 934-936 [*ZhETF* **56** (1969) 1742-1747 (in Russian)].
- [376] M.P. Ryan and L.C. Shepley, *Homogeneous relativistic cosmologies* (Princeton Univ. Press: Princeton, 1975).
- [377] R. Sachs, *Gravitational waves in general relativity. IV. The outgoing radiation condition*, *Proc. Roy. Soc.* **A264** (1961) 309-338.
- [378] R.K. Sachs and A.M. Wolfe, *Perturbations of a cosmological model and angular variations of the microwave background*, *Astrophys. J.* **147** (1967) 73-90.
- [379] P.D. Sackett, *Inflationary model predicts little rotation: cosmology*, *Science News* **124** (1983) 20.
- [380] R.X. Saibatalov, *Electromagnetic field in causal and acausal Gödel-type space-times*, *Gen. Rel. Grav.* **27** (1995) 697-711.
- [381] M. Sakoto, *Closed conformal infinitesimal transformations of Tolman-Omer and Gödel cosmological spaces*, *Comptes Rendus Acad. Sci. (Paris), Ser. A: Sci. Math.* **284** (1977) 633-636 (in French).
- [382] J.L. Sanz, *Dynamical importance of vorticity and shear in the universe*, *J. Math. Phys.* **23** (1982) 1732-1736.
- [383] M. Scherfner, *History of cosmic rotation and new developments*, in: “Current topics in mathematical cosmology”, *Proc. of Intern. Seminar, Potsdam, 30 March-4 April 1998*, Eds. M. Rainer and H.-J. Treder (Singapore: World Scientific, 1998) 419-424.
- [384] A.A. Schmidt, C.A.Z. Vasconcellos, and H.A. Dottori, *Cosmological mass density and*

- the Regge law*, *Rev. Mexicana Astro. Astrof.* **14** (1987) 66-70.
- [385] E. Schmutzer, *Relativistische Physik. Klassische Theorie* (Akad. Verlag: Leipzig, 1958).
- [386] E. Schücking, *Homogene scherungsfreie Weltmodelle in der relativistischen Kosmologie*, *Naturwiss.* **19** (1957) 507.
- [387] E. Schücking and O. Heckmann, *World models*, in: *La structure et l'évolution de l'Universe*, Ed. R. Stoops (Bruxelles, 1958) 149-159.
- [388] D. Sciama, *Issues in cosmology*, in: "Some strangeness in proportion: A Centennial Symp. of A. Einstein", Ed. H. Woolf (Addison-Wesley: NY, 1980) 387-404.
- [389] S. Seitz, P. Schneider, and J. Ehlers, *Light propagation in arbitrary spacetimes and the gravitational lens approximation*, *Class. Quantum Grav.* **11** (1994) 2345-2373.
- [390] G.N. Shikin, *On the influence of electromagnetic and real scalar fields on a complex massive scalar field in the Gödel universe*, in: "Probl. Theor. Grav. and Elem. Part.", Ed. K.P. Stanyukovich (Moscow: Energoatomizdat, 1985) **15**, 98-102 (in Russian).
- [391] J. Silk, *Local irregularities in a Gödel universe*, *Astrophys. J.* **143** (1966) 689-699.
- [392] J. Silk, *The instability of a rotating universe*, *Mon. Not. R. Astr. Soc.* **147** (1970) 13-19.
- [393] J. Silk, *Primordial turbulence and intergalactic matter*, in: *Cargèse Lect. in Phys.* vol. **6**, Ed. E. Schatzman (Gordon and Breach: NY, 1973) 401-503.
- [394] S. Singh, *Cosmological models with shear and rotation*, *J. Astrophys. and Astron.* **20** (1999) 67-77.
- [395] T. Singh and A.K. Agrawal, *Stationary universe in general relativity and Lyra's geometry*, *Fortsch. Phys.* **42** (1994) 71-93.
- [396] N.I. Singh, *Slow rotational perturbations of Robertson-Walker universes*, *Astrophys.*

- and *Space Sci.* **148** (1988) 199-205.
- [397] N.I. Singh and C.G. Singh, *Slowly rotating cosmological universe filled with viscous fluid*, *Astrophys. and Space Sci.* **253** (1997) 189-194.
- [398] R. Sistero, *Rotation of the universe and the universal angular momentum – mass relationship*, *Astrophys. Lett.* **23** (1983) 235-237.
- [399] G.V. Skrotskii, *The influence of gravitation on the propagation of light*, *Sov. Phys. Doklady* **2** (1957) 226-229.
- [400] L.L. Smalley, *Gödel cosmology in Riemann-Cartan space-time with spin density*, *Phys. Rev.* **D32** (1985) 3124-3127.
- [401] L.L. Smalley, *Self-consistent Gödel cosmology with spin density in Riemann-Cartan space-time*, *Phys. Lett.* **A113** (1986) 463-466.
- [402] G.F. Smoot et al, *Structure in the COBE Differential Microwave Radiometer first year maps*, *Astrophys. J.* **396** (1992) L1-5.
- [403] H.H. Soleng, *Does cosmos really have a fast rotation?*, *Ann. d. Phys. (DDR)* **45** (1988) 507-508.
- [404] H.H. Soleng, *Gödel-type cosmologies with spin density*, *Phys. Lett.* **A137** (1989) 326-328.
- [405] M.M. Som and A.K. Raychaudhuri, *Cylindrically symmetric charged dust distributions in rigid rotation in general relativity*, *Proc. Roy. Soc. (Lond.)* **A304** (1968) 81-86.
- [406] H. Stein, *On the paradoxical time-structures of Gödel*, *Philosophy of Science* **37** (1970) 589-601.
- [407] W.J. van Stockum, *The gravitational field of a distribution of particles rotating about an axis of symmetry*, *Proc. Roy. Soc. Edinburgh* **57** (1936/37) 135-154.

- [408] W.R. Stoeger, M.E. Araujo, and T. Gebbie, *The limits on cosmological anisotropies and inhomogeneities from COBE data*, *Astrophys. J.* **476** (1997) 435-439.
- [409] N. Straumann, *General relativity and relativistic astrophysics*, *Texts and Monographs In Physics* (Springer: Berlin, 1984).
- [410] G.C. Surpi, and D.D. Harari, *Weak lensing by large-scale structure and the polarization properties of distant radio-sources*, *Astrophys. J.* **515** (1999) 455-464.
- [411] E. Sviestins, *Null geodesics, caustics and apparent motion of galaxies in a finite rotating universe*, *Gen. Rel. Grav.* **16** (1984) 161-173.
- [412] E. Sviestins, *Some rotating, time-dependent Bianchi type-IX cosmologies with heat flow*, *Gen. Rel. Grav.* **17** (1985) 521-523.
- [413] L.J. Tassie, *Cosmic strings, superstrings and the evolution of the universe*, *Nature* **323** (1986) 40-42.
- [414] A.F. Teixeira, M. Rebouças, and J. Åman, *Isometries of homogeneous Gödel-type spacetimes*, *Phys. Rev.* **D32** (1985) 3309-3311.
- [415] K. Tomita, *Theoretical relations among observable quantities in an anisotropic and homogeneous universe*, *Progr. Theor. Phys.* **40** (1968) 264-276.
- [416] A. Trautman, *Spin and torsion may avert gravitational singularity*, *Nature Phys. Sci.* **242** (1973) 7-8.
- [417] H.-J. Treder, *Aberrations-Konstante und Rotation des Kosmos*, *Ann. d. Phys.* **42** (1985) 71-72.
- [418] H.-J. Treder, *Das rotierende Universum*, *Wissenschaft und Fortschritt*, **37** (1987) 100-102.
- [419] A.Yu. Ushakov and A.D. Chernin, *The structure of protogalactic eddies*, *Sov. Astron.* **27** (1983) 367-370 [*Astron. Zh* **60** (1983) 625-631 (in Russian)].

- [420] J.K. Usham and R.K.T. Singh, *Slowly rotating cosmological viscous fluid universe in Brans-Dicke theory*, *Astrophys. and Space Sci.* **189** (1992) 79-87.
- [421] P.C. Vaidya, *An Einstein-Gödel universe*, *Gen. Rel. Grav.* **9** (1978) 801-807.
- [422] P.C. Vaidya and L.K. Patel, *A rotating homogeneous universe with an electromagnetic field*, *Gen. Rel. Grav.* **16** (1984) 355-364.
- [423] E.P.V. Vaidya, M.L. Bedran, and M.M. Som, *Comments on the source of Gödel type metric*, *Progr. Theor. Phys.* **72** (1984) 757-759.
- [424] F. Valdes, J.A. Tyson, and J.F. Jarvis, *Alignment of faint galaxy images: cosmological distortion and rotation*, *Astron. J.* **271** (1983) 431-441.
- [425] V.M. Villalba, *Dirac spinor in a nonstationary Gödel-type cosmological universe*, *Mod. Phys. Lett.* **A8** (1993) 3011-3018.
- [426] A.M. Volkov, A.A. Izmet'sev, and G.V. Skrotskii, *The propagation of electromagnetic waves in a Riemannian space*, *Sov. Phys. JETP* **32** (1971) 686-689 [*ZhETF* **59** (1970) 1254-1261 (in Russian)].
- [427] M. Walker and R. Penrose, *On quadratic first integrals of the geodesic equations for type {22} spacetimes*, *Commun. Math. Phys.* **18** (1970) 265-274.
- [428] J.F.C. Wardle, R.A. Perley, and M.H. Cohen, *Observational evidence against birefringence over cosmological distances*, *Phys. Rev. Lett.* **79** (1997) 1801-1804.
- [429] N.P. Warner and H.A. Buchdahl, *On the world function of the Gödel metric*, *J. Phys.: Math. and Gen.* **A13** (1980) 509-516.
- [430] C.F. von Weizsäcker, *Zur Kosmogonie*, *Z. Astrophysik* **24** (1948) 181-206.
- [431] P.S. Wesson, *Discrete source counts and the rotation of the local universe in hierarchical cosmology*, *Astrophys. and Space Sci.* **37** (1975) 235-256.

- [432] E.T. Whittaker, *Spin in the universe*, Yearbook of Roy. Soc. Edinburgh (1945) 5-13.
- [433] J.G. Williams, $(2 + 1)$ -dimensional spacetime with rotation and expansion, *Gen. Rel. Grav.* **26** (1994) 499-512.
- [434] J.G. Williams, *Thermodynamics of a rotating and expanding spacetime in 2+1 dimensions*, in: “Proc. of the 5th Canadian Conf. on General Relativity and Relativistic Astrophysics, Waterloo, 13-15 May 1993”. Eds. R.B.Mann and R.G. McLenaghan (World Scientific: Singapore, 1994), 477-480.
- [435] A.M. Wolfe, *New limits on the shear and rotation of the universe from the X-ray background*, *Astrophys. J.* **159** (1970) L61-67.
- [436] E.L. Wright et al, *Interpretation of the cosmic microwave background radiation anisotropy detected by the COBE Differential Microwave Radiometer*, *Astrophys. J.* **396** (1992) L13-18.
- [437] J.P. Wright, *Solution of Einstein’s field equations for a rotating, stationary, and dust-filled universe*, *J. Math. Phys.* **6** (1965) 103-105.
- [438] I. Yavuz, C. Battal, and H. Baysal, *Slow rotational perturbations of Friedmann universes*, *Astrophys. Space Sci.* **167** (1990) 341-345.
- [439] I. Yavuz and H. Baysal, *Nonstatic Gödel-type cosmological model with perfect fluid and heat flow*, *Int. J. Theor. Phys.* **33** (1994) 2285-2289.
- [440] L. Zhen, D. Gan, and C. Lin, *Characteristic actions $\hbar^{(s)}$ in the structure of the universe*, *Astrophys. Space Sci.* **116** (1986) 215-224.